

**Eight Repeatedly Documented Findings Each Show that EMF Safety Guidelines Do Not Predict Biological Effects and Are, Therefore Fraudulent:  
The Consequences for Both Microwave Frequency Exposures and Also 5G  
3rd Edition**

Martin L. Pall, PhD, Professor Emeritus of Biochemistry and Basic Medical Sciences, Washington State University, [martin\\_pall@wsu.edu](mailto:martin_pall@wsu.edu)

**Abstract**

ICNIRP, US FCC, EU and other EMF safety guidelines are all based on the assumption that average EMF intensities and average SAR can be used to predict biological effects and therefore safety. Eight different types of quantitative or qualitative data are analyzed here to determine whether these safety guidelines predict biological effects. In each case the safety guidelines fail and in most of these, fail massively. Effects occur at approximately 100,000 times below allowable levels and the basic structure of the safety guidelines is shown to be deeply flawed. The safety guidelines ignore demonstrated biological heterogeneity and established biological mechanisms. Even the physics underlying the safety guidelines is shown to be flawed. Pulsed EMFs are in most cases much more biologically active than are non-pulsed EMFs of the same average intensity, but pulsations are ignored in the safety guidelines despite the fact that almost all of our current exposures are highly pulsed. There are exposure windows such that maximum effects are produced in certain intensity windows and also in certain frequency windows but the consequent very complex dose-response curves are ignored by the safety guidelines. Several additional flaws in the safety guidelines are shown through studies of both individual and paired nanosecond pulses. The properties of 5G predict that guidelines will be even more flawed in predicting 5G effects than the already stunning flaws that the safety guidelines have in predicting our other EMF exposures. The consequences of these findings is that “safety guidelines” should always be expressed in quotation marks; they do not predict biological effects and therefore do not predict safety. Because of that we have a multi-trillion dollar set of companies, the telecommunication industry, where all assurances of safety are fraudulent because they are based on these “safety guidelines.”

Introduction

The current safety guidelines including the EU safety guideline, the US FCC safety guidelines, the 2013 UK safety guidelines, Canada’s safety code 6 and the recent revisions proposed in the 2018 ICNIRP draft, are all very similar to although modified from of the ICNIRP 1998 safety guidelines. The viability of each of these is dependent on the viability of the 1998 ICNIRP safety guidelines [International Commission on non-ionizing radiation protection. 1998 ICNIRP GUIDELINES FOR LIMITING EXPOSURE TO TIME-VARYING ELECTRIC, MAGNETIC AND ELECTROMAGNETIC FIELDS (UP TO 300 GHZ) Health Physics 74 (4):494-522]. There are ongoing processes to make the “safety guidelines” even looser to allow the rollout of 5G and while these are important, they are not considered in this paper.

**Table 1: 1998 ICNIRP “Safety Guidelines”**

	<b>Frequency range</b>	<b>Whole-body SAR (averaged over 6 minutes)</b>	<b>Localized SAR (head &amp; trunk, averaged over 6 minutes)</b>	<b>Localized SAR (limbs, averaged over 6 minutes)</b>
Occupational	100 KHz-	0.4 (W/kg)	10 (W/kg)	20 (W/kg)

exposure	10 GHz			
General public exposure	100 KHz-10 GHz	0.08 (W/kg)	2 (W/kg)	4 (W/kg)

There are three points that need to be considered:

- Because specific absorption rates (SAR) only predict thermal (heating) effects, there is no reason to assume that these “safety guidelines” predict non-thermal effects.
- There is no reason why effects that occur with very brief exposures should be assumed to be predicted by average exposures over 6 minutes. Some of the more recent guidelines, including the recent ICNIRP draft and the FCC occupational exposure guidelines are averaged over 30 minutes, making this issue of averaging still more problematic.
- When one is concerned about non-thermal responses that are localized, such as local oxidative stress or apoptosis or local DNA effects, there is no reason to use much less stringent guidelines for localized exposures as compared with whole body exposures.

These “safety guidelines” are the basis of the guidelines from the EU, the US FCC, Canada’s safety code 6 and others and while there are some minor differences, these ICNIRP levels can be taken as being similar to each of them. Each of these use exposures averaged over 6 minutes or 30 minutes, where allowable levels are based on SAR and, therefore, only protect us from thermal effects. For example, the EU 1999 general public safety guidelines, COUNCIL RECOMMENDATION of 12 July 1999 on the limitation of exposure of the general public to electromagnetic fields (100 KHz to 300 GHz) are identical to those listed by ICNIRP in Table 1 (<https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:31999H0519&from=EN>) and these same general public guidelines were repeated by the EU in 2013 (L 79, 29.6.2013 Official Journal of the European Union). Similarly, the The Australian (ARPANSA) April, 2002 “safety guidelines” for both general public and occupational exposures are identical to the 1998 ICNIRP guidelines (<https://www.arpansa.gov.au/sites/default/files/legacy/pubs/rps/rps3.pdf>).

The 1999 FCC “safety guidelines” are listed in Table 2, below and the similarities to and also some differences with the ICNIRP guidelines, shown in Table 1 can be seen.

**Table 2: 1999 US FCC occupational and general public limits for both whole body and localized (partial body) exposures**

	Frequency range	Whole body SAR	Localized (partial body) SAR
Occupational exposure	100 KHz – 6 GHz	<0.4 W/kg averaged over 30 minutes	<8 W/kg averaged over 30 minutes
General public exposure	100 KHz – 6 GHz	<0.08 W/kg averaged over 6 minutes	<1.6 W/kg averaged over 6 minutes

The same three concerns expressed above regarding the ICNIRP guidelines are also concerns with respect to the FCC guidelines. The occupational FCC guidelines are weaker than the 1998 ICNIRP guidelines because it uses 30 minute as opposed to 6 minute averaging. The general public whole body guidelines, possibly the most important, are identical for the two guidelines. The localized (partial body) FCC levels are slightly more stringent than are 1998 ICNIRP levels. There are some differences, but the overall structure of both is very similar, with both based on average intensities or SAR and both, therefore, only possibly predicting thermal effects.

We have, then three points that were raised, above, each of which raise serious questions about the “safety-guidelines.” But no information is provided to this point in this paper, as to whether these guidelines predict biological effects and therefore safety or not. This paper is mainly focused on the following question: How then do the allowable “safety guideline” exposure levels compare with levels found in empirical studies to produce actual effects? Eight distinct types of repeatedly found patterns of evidence are considered here, each of which clearly show that the “safety guidelines” do not predict biological effects.

1. I list here bodies of evidence from published reviews, that clearly show that non-thermal exposures to microwave and other frequency EMF, at levels far below “safety guideline” allowable levels, produce each of 9 different types of important health-related effects. Many of the citations listed here are from my 90 page EMF document but substantial numbers of new findings are listed here, as well.

These effects are as follows:

1) **Lowered fertility, including tissue remodeling changes in the testis, lowered sperm count and lowered motility and other measures of lowered sperm quality, lowered female fertility including ovarian remodeling, oocyte (follicle) loss, lowered estrogen, progesterone and testosterone levels (that is sex hormone levels), increased spontaneous abortion incidence, lowered libido** (33 reviews).

1. Glaser ZR, PhD. 1971 Naval Medical Research Institute Research Report, June 1971. Bibliography of Reported Biological Phenomena (“Effects”) and Clinical Manifestations Attributed to Microwave and Radio-Frequency Radiation. Report No. 2 Revised.  
[https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as\\_sdt=0%2C38](https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as_sdt=0%2C38) (Accessed Sept. 9, 2017)
2. Tolgskaya MS, Gordon ZV. 1973. Pathological Effects of Radio Waves, Translated from Russian by B Haigh. Consultants Bureau, New York/London, 146 pages.
3. Leach WM. 1980 Genetic, growth and reproductive effects of microwave radiation. Bull N Y Acad Med 56:249-257.
4. Goldsmith JR. 1997 Epidemiological evidence relevant to radar (microwave) effects. Environ Health Perspect 105(Suppl 6):1579-1587.
5. Aitken RJ, De Iuliis GN. 2007 Origins and consequences of DNA damage in male germ cells. Reprod Biomed Online 14:727-733.
6. Hazout A, Menezo Y, Madelenat P, Yazbeck C, Selva J, Cohen-Bacrie P. 2008 [Causes and clinical implications of sperm DNA damages]. Gynecol Obstet Fertil ;36:1109-1117.
7. Makker K, Varghese A, Desai NR, Mouradi R, Agarwal A. 2009 Cell phones: modern man's nemesis? Reprod Biomed Online 18:148-157.
8. Panagopoulos DJ, Margaritis LH. 2009 Biological and Health Effects of Mobile Telephone Radiations. Int J Med Biol Front 2009; 15 (1-2): 33-76.
9. Kang N, Shang XJ, Huang YF. 2010 [Impact of cell phone radiation on male reproduction]. Zhonghua Nan Ke Xue 16:1027-1030.
10. Gye MC, Park CJ. 2012 Effect of electromagnetic field exposure on the reproductive system. Clin Exp Reprod Med 39:1-9. doi.org/10.5653/cerm.2012.39.1.1

11. La Vignera S, Condorelli RA, Vicari E, D'Agata R, Calogero AE. 2012 Effects of the exposure to mobile phones on male reproduction: a review of the literature. *J Androl* 33:350-356.
12. Carpenter DO. 2013 Human disease resulting from exposure to electromagnetic fields. *Rev Environ Health* 28:159-172.
13. Nazıroğlu M, Yüksel M, Köse SA, Özkaya MO. 2013 Recent reports of Wi-Fi and mobile phone-induced radiation on oxidative stress and reproductive signaling pathways in females and males. *J Membr Biol* 246:869-875.
14. Adams JA, Galloway TS, Mondal D, Esteves SC, Mathews F. 2014 Effect of mobile telephones on sperm quality: a systematic review and meta-analysis. *Environ Int* 70:106-112.
15. Hasan GM, Sheikh IA, Karim S, Haque A, Kamal MA, Chaudhary AG, Azhar E, Mirza Z. 2014 Effect of electromagnetic radiations on neurodegenerative diseases- technological revolution as a curse in disguise. *CNS Neurol Disord Drug Targets* 13:1406-1412.
16. Liu K, Li Y, Zhang G, Liu J, Cao J, Ao L, Zhang S. 2014 Association between mobile phone use and semen quality: a systematic review and meta-analysis. *Andrology* 2:491-501.
17. K Sri N. 2015 Mobile phone radiation: physiological & pathophysiological considerations. *Indian J Physiol Pharmacol* 59:125-135.
18. Hensinger P, Wilke E. 2016. Mobilfunk-Studienergebnisse bestätigen Risiken Studienrecherche 2016-4 veröffentlicht. *Umwelt Medizin Gesellschaft* 29:3/2016.
19. Starkey S. 2016 Inaccurate official assessment of radiofrequency safety by the Advisory Group on Non-ionising Radiation. *Rev Environ Health* 31:493-503.
20. Houston BJ, Nixon B, King BV, De Iuliis GN, Aitken RJ. 2016 The effects of radiofrequency electromagnetic radiation on sperm function. *Reproduction* 152:R263-R276
21. Pall ML. 2018 Wi-Fi is an important threat to human health. *Environ Res* 164:404-416.
22. Kocaman A, Altun G, Kaplan AA, Deniz OG, Yurt KK, Kaplan S. 2018 Genotoxic and carcinogenic effects of non-ionizing electromagnetic fields. *Environ Res* 163: 71-79. doi: 10.1016/j.envres.2018.01.034
23. Yahyazadeh A, Deniz ÖG, Kaplan AA, Altun G, Yurt KK, Davis D. 2018 The genomic effects of cell phone exposure on the reproductive system. *Environ Res*. 2018 Nov;167:684-693. doi: 10.1016/j.envres.2018.05.017.
24. Belpomme D, Hardell L, Belyaev I, Burgio E, Carpenter DO. 2018 Thermal and non-thermal health effects of low intensity non-ionizing radiation: An international perspective. *Environ Pollut* 2018 Nov;242(Pt A):643-658. doi: 10.1016/j.envpol.2018.07.019.
25. Wilke I. 2018 Biological and pathological effects of 2.45 GHz on cells, fertility, brain and behavior. *Umwelt Medizin Gessellschaft* 2018 Feb 31 (1).
26. Kesari KK, Agarwal A, Henkel R. Radiations and male fertility. 2018 *Reprod Biol Endocrinol*. 2018 Dec 9;16(1):118. doi: 10.1186/s12958-018-0431-1.
27. Belyaev I. 2018 Health effects of chronic exposure to radiation from mobile communication. Chapter 4 in: *Mobile Communications and Public Health*, Marko Markov, Ed., CRC Press, pp 65-99.
28. Santini SJ, Cordone V, Falone S, Mijit M, Tatone C, Amicarelli F, Di Emidio G. 2018 Role of Mitochondria in the Oxidative Stress Induced by Electromagnetic Fields: Focus on Reproductive Systems. *Oxid Med Cell Longev*. 2018 Nov 8;2018:5076271. doi: 10.1155/2018/5076271.

29. Zha X-D, Wang W-W, Xu S, Shang X-J. 2019. [Impacts of electromagnetic radiation from cell phones and Wi-Fi on spermatogenesis]. *Zhonghua Nan Ke Xue* 25:451-455.
30. Jaffar FHJ, Osman K, Ismail NH, Chn K-Y. 2019. Adverse Effects of Wi-Fi Radiation on Male Reproductive System: A Systematic Review. *Tohoku J Exp Med* 248:169-179. doi: 10.1620/tjem.248.169.
31. Yadav H, Rai U., Singh R. 2021. Radiofrequency radiation: A possible threat to male fertility. *Reprod Toxicol.* 100:90-100.
32. Negi, P., Singh, R. 2021. Association between reproductive health and nonionizing radiation exposure. *Electromagn Biol Med.* 40:92-102.
33. Uche U. 2022. How exposure to cell phones and other technology affects reproductive health (Environmental Working Group). <https://www.ewg.org/research/how-exposure-cell-phones-and-other-technology-affects-reproductive-health>

- 2) **Neurological/neuropsychiatric effects including EEG and transmitter function; widespread neurological dysfunction, including: sleep disturbance/insomnia; fatigue/tiredness; headache; depression/depressive symptoms; lack of concentration/attention/cognitive dysfunction; dizziness/vertigo; memory changes; restlessness/tension/anxiety/stress/agitation; irritability; tinnitus** (44 reviews). It should not be surprising that there are so many brain related effects, because the Tolgskaya and Gorden review (#3 below) finds that in rodents the nervous system including the brain is the most sensitive organ in the body to histological change produced by non-thermal EMF exposures, followed by the heart and the testis.
1. Marha K. 1966 Biological Effects of High-Frequency Electromagnetic Fields (Translation). ATD Report 66-92. July 13, 1966 (ATD Work Assignment No. 78, Task 11). <http://www.dtic.mil/docs/citations/AD0642029> (accessed March 12, 2018)
  2. Glaser ZR, PhD. 1971 Naval Medical Research Institute Research Report, June 1971. Bibliography of Reported Biological Phenomena (“Effects”) and Clinical Manifestations Attributed to Microwave and Radio-Frequency Radiation. Report No. 2 Revised. [https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as\\_sdt=0%2C38](https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as_sdt=0%2C38) (Accessed Sept. 9, 2017)
  3. Tolgskaya MS, Gordon ZV. 1973. Pathological Effects of Radio Waves, Translated from Russian by B. Haigh. Consultants Bureau, New York/London, 146 pages.
  4. Bise W. 1978 Low power radio-frequency and microwave effects on human electroencephalogram and behavior. *Physiol Chem Phys* 10:387-398.
  5. Raines, J. K. 1981. Electromagnetic Field Interactions with the Human Body: Observed Effects and Theories. Greenbelt, Maryland: National Aeronautics and Space Administration 1981; 116 p.
  6. Frey AH. 1993 Electromagnetic field interactions with biological systems. *FASEB J* 7:272-281.
  7. Lai H. 1994 Neurological effects of radiofrequency electromagnetic radiation. In: *Advances in Electromagnetic Fields in Living Systems*, Vol. 1, J.C. Lin, Ed., Plenum Press, New York, pp. 27-88.
  8. Grigor'ev IuG. 1996 [Role of modulation in biological effects of electromagnetic radiation]. *Radiats Biol Radioecol* 36:659-670.

9. Lai, H 1998 Neurological effects of radiofrequency electromagnetic radiation. [http://www.mapcruzin.com/radiofrequency/henry\\_lai2.htm](http://www.mapcruzin.com/radiofrequency/henry_lai2.htm).
10. Stepansky R, Jahn O, Windischbauer G, Zeitlhofer J. 2000 [Electromagnetic fields--effects on health]. *Acta Med Austriaca*. 2000;27(3):69-77.
11. Lebedeva NN, Sulimov AV, Sulimova OP, Korotkovskaya TI, Gailsu T. 2000 Investigations of brain potentials in humans exposed to the electromagnetic field of mobile phones. *Crit Rev Biomed Engineer* 28:323-337.
12. Hecht K. 2001. [Effects of electromagnetic fields: A review of Russian study results 1960-1996]. *Umwelt-Medizin-Gesellschaft* 14: 222-231.
13. Westerman R, Hocking B. 2004 Diseases of modern living: neurological changes associated with mobile phones and radiofrequency radiation in humans. *Neurosci Lett* 361:13-16.
14. Oberfeld, G., Navarro, A.E., Portoles, M., Maestu, C., Gomez-Perretta, C., 2004. The microwave syndrome: further aspects of a Spanish study, <http://www.apdr.info/electrocontaminacion/Documentos/Investigacion/ESTUDOS%20EPIDEMIOLOGICOS%20E%20ANTENAS/The%20Microwave%20Syndrome%20-%20Further%20Aspects%20of%20a%20Spanish%20Study.pdf>.
15. **Abdel-Rassoul G, El-Fateh OA, Salem MA, Michael A, Farahat F, El-Batanouny M, Salem E. 2007. Neurobehavioral effects among inhabitants around mobile phone base stations. *Neurotoxicology* 28(2):434-40. doi: 10.1016/j.neuro.2006.07.012.**
16. **Orendacova J, Orendac M, Racekova E, Marsala L. 2007. Neurobiological effects of microwave exposure: A review focused on morphological findings in experimental animals. *Arch Ital de Biologie* 145:1-12.**
17. Hardell, L., Sage, C. 2008. Biological effects from electromagnetic field exposure and public exposure standards. *Biomed. Pharmacother.* 62, 104-109.
18. Makker K, Varghese A, Desai NR, Mouradi R, Agarwal A. 2009 Cell phones: modern man's nemesis? *Reprod Biomed Online* 18:148-157.
19. **Khurana VG, Hardell L, Everaert J, Bortkiewicz A, Carlberg M, Ahonen M. 2010 Epidemiological evidence for a health risk from mobile phone base stations. *Int J Occup Environ Health* 16:263-267.**
20. **Levitt, B. B., Lai, H. 2010. Biological effects from exposure to electromagnetic radiation emitted by cell tower base stations and other antenna arrays. *Environ. Rev.* 18, 369-395. doi.org/10.1139/A10-018**
21. Bak M, Zmysłony M. 2010 [Effects of electromagnetic field from cellular phones on selected central nervous system functions: a literature review]. *Med Pr.* 2010;61(6):671-83.
22. **Bortkiewicz, A., Gadzicka, E., Szyjkowska, A., Politański, P., Mamrot, P., Szymczak, W., Zmysłony, M., 2012. Subjective complaints of people living near mobile phone base stations in Poland. *Int. J. Occup. Med. Environ. Health* 25, 31-40.**
23. Carpenter DO. 2013 Human disease resulting from exposure to electromagnetic fields. *Rev Environ Health* 2013;28:159-172.
24. Havas M. 2013 Radiation from wireless technology affects the blood, the heart and the autonomic nervous system. *Rev Environ Health* 28: 75-84.
25. Kesari KK, Siddiqui MH, Meena R, Verma HN, Kumar S. 2013. Cell phone radiation exposure on brain and associated biological systems. *Ind J Exp Biol* 51: 187-200.
26. Hasan GM, Sheikh IA, Karim S, Haque A, Kamal MA, Chaudhary AG, Azhar E, Mirza Z. 2014 Effect of electromagnetic radiations on neurodegenerative diseases- technological revolution as a curse in disguise. *CNS Neurol Disord Drug Targets* 13:1406-1412.

27. Sage C. 2015 The implications of radiofrequency safety by the Advisory Group on Non-ionising Radiation. *Rev Environ Health* 31:493-503.
28. Bhargav H, Srinivasan TM, Varambally S, Gangadhar BN, Koka P. 2015 Effect of Mobile Phone-Induced Electromagnetic Field on Brain Hemodynamics and Human Stem Cell Functioning: Possible Mechanistic Link to Cancer Risk and Early Diagnostic Value of Electronphotonic Imaging. *J Stem Cells*. 2015;10(4):287-94.
29. Politański P, Bortkiewicz A, Zmysłony M. 2016 [Effects of radio- and microwaves emitted by wireless communication devices on the functions of the nervous system selected elements]. *Med Pr* 67:411-421.
30. Hensinger P, Wilke E. 2016. Mobilfunk-Studienergebnisse bestätigen Risiken Studienrecherche 2016-4 veröffentlicht. *Umwelt Medizin Gesellschaft* 29:3/2016.
31. Pall ML. 2016 Microwave frequency electromagnetic fields (EMFs) produce widespread neuropsychiatric effects including depression. *J Chem Neuroanat* 75(Pt B):43-51. doi: 10.1016/j.jchemneu.2015.08.001.
32. Hecht, Karl. 2016 Health Implications of Long-Term Exposures to Electrosmog. Brochure 6 of A Brochure Series of the Competence Initiative for the Protection of Humanity, the Environment and Democracy. [http://kompetenzinitiative.net/KIT/wp-content/uploads/2016/07/KI\\_Brochure-6\\_K\\_Hecht\\_web.pdf](http://kompetenzinitiative.net/KIT/wp-content/uploads/2016/07/KI_Brochure-6_K_Hecht_web.pdf) (accessed Feb. 11, 2018)
33. Starkey S. 2016 Inaccurate official assessment of radiofrequency safety by the Advisory Group on Non-ionising Radiation. *Rev Environ Health* 31:493-503.
34. Sangün Ö, Dündar B, Çömlekçi S, Büyükgebiz A. 2016 The Effects of Electromagnetic Field on the Endocrine System in Children and Adolescents. *Pediatr Endocrinol Rev* 13:531-545.
35. Belyaev I, Dean A, Eger H, Hubmann G, Jandrisovits R, Kern M, Kundi M, Moshammer H, Lercher P, Müller K, Oberfeld G, Ohnsorge P, Pelzmann P, Scheingraber C, Thill R. 2016 EUROPAEM EMF Guideline 2016 for the prevention, diagnosis and treatment of EMF-related health problems and illnesses. *Rev Environ Health* DOI 10.1515/reveh-2016-0011.
36. Zhang J, Sumich A, Wang GY. 2017 Acute effects of radiofrequency electromagnetic field emitted by mobile phone on brain function. *Bioelectromagnetics* 38:329-338. doi: 10.1002/bem.22052.
37. Belpomme D, Hardell L, Belyaev I, Burgio E, Carpenter DO. 2018 Thermal and non-thermal health effects of low intensity non-ionizing radiation: An international perspective. *Environ Pollut* 2018 Nov;242(Pt A):643-658. doi: 10.1016/j.envpol.2018.07.019.
38. Lai H. 2018. A Summary of Recent Literature (2007–2017) on Neurological Effects of Radio Frequency Radiation. Chapter 8 in *Mobile Communications and Public Health*, Marko Markov, Ed., CRC press, pp 189-224.
39. Belyaev I. 2018 Health effects of chronic exposure to radiation from mobile communication. Chapter 4 in: *Mobile Communications and Public Health*, Marko Markov, Ed., CRC Press, pp 65-99.
40. Pall ML. 2018 Wi-Fi is an important threat to human health. *Environ Res* 164:404-416.
- 41. Subhan F, Khan A, Ahmed S, Malik SN, Bakshah ST, Tahir S.. 2018 Mobile antenna's and its impact on human health. *J Medical Imaging and Health Informatics*. 8(66):11266 - 1273. doi: 10.1166//jjmihi.22018.2296.**
42. Wilke I. 2018 Biological and pathological effects of 2.45 GHz on cells, fertility, brain and behavior. *Umwelt Medizin Gesselshaft* 2018 Feb 31 (1).
43. Narayanan SN, Jetti R, Kesari KK, Kumar RS, Nayak SB, Bhat PG. 2019 Radiofrequency electromagnetic radiation-induced behavioral changes and their possible basis. *Environ Sci Pollut Res Int*. 2019 Oct;26(30):30693-30710. doi: 10.1007/s11356-019-06278-5.

44. Isaković J1, Gorup D, Mitrečić D. 2019. Molecular mechanisms of microglia- and astrocyte-driven neurorestoration triggered by application of electromagnetic fields. *Croatian Med J* 60:127-140. doi: 10.3325/cmj.2019.60.127
- 3) **Effects on cellular DNA including single strand and double strand breaks in cellular DNA and on oxidized bases in cellular DNA; also evidence for chromosomal mutations produced by double strand DNA breaks. These produce all of the important type of mutations, as described at the DNA level that have roles in cancer causation and in human whole organism mutation** (27 reviews).
1. Glaser ZR, PhD. 1971 Naval Medical Research Institute Research Report, June 1971. Bibliography of Reported Biological Phenomena (“Effects”) and Clinical Manifestations Attributed to Microwave and Radio-Frequency Radiation. Report No. 2 Revised. [https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as\\_sdt=0%2C38](https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as_sdt=0%2C38) (Accessed Sept. 9, 2017)
  2. Leach WM. 1980 Genetic, growth and reproductive effects of microwave radiation. *Bull N Y Acad Med* 56:249-257.
  3. Goldsmith JR. 1997 Epidemiologic evidence relevant to radar (microwave) effects. *Environ Health Perspect* 105(Suppl 6):1579-1587.
  4. Yakymenko IL, Sidorik EP, Tsybulin AS. 1999 [Metabolic changes in cells under electromagnetic radiation of mobile communication systems]. *Ukr Biokhim Zh* (1999), 2011 Mar-Apr:20-28.
  5. Aitken RJ, De Iuliis GN. 2007 Origins and consequences of DNA damage in male germ cells. *Reprod Biomed Online* 14:727-733.
  6. Hardell, L., Sage, C. 2008. Biological effects from electromagnetic field exposure and public exposure standards. *Biomed. Pharmacother.* 62, 104-109.
  7. Hazout A, Menezo Y, Madelenat P, Yazbeck C, Selva J, Cohen-Bacrie P. 2008 [Causes and clinical implications of sperm DNA damages]. *Gynecol Obstet Fertil* ;36:1109-1117.
  8. Phillips JL, Singh NP, Lai H. 2009 Electromagnetic fields and DNA damage. *Pathophysiology* 16:79-88.
  9. Panagopoulos DJ, Margaritis LH. 2009 Biological and Health Effects of Mobile Telephone Radiations. *Int J Med Biol Front* 2009; 15 (1-2): 33-76.
  10. Ruediger HW. 2009 Genotoxic effects of radiofrequency electromagnetic fields. *Pathophysiology.* 16:89-102.
  11. Makker K, Varghese A, Desai NR, Mouradi R, Agarwal A. 2009 Cell phones: modern man's nemesis? *Reprod Biomed Online* 18:148-157.
  12. Yakymenko I, Sidorik E. 2010 Risks of carcinogenesis from electromagnetic radiation and mobile telephony devices. *Exp Oncol* 32:729-736.
  13. Yakymenko IL, Sidorik EP, Tsybulin AS. 2011 [Metabolic changes in cells under electromagnetic radiation of mobile communication systems]. *Ukr Biokhim Zh* (1999). 2011 Mar-Apr;83(2):20-28.
  14. Gye MC, Park CJ. 2012 Effect of electromagnetic field exposure on the reproductive system. *Clin Exp Reprod Med* 39:1-9. doi.org/10.5653/cerm.2012.39.1.1
  15. Pall, ML. 2013. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med* 17:958-965. doi: 10.1111/jcmm.12088.



16. Hasan GM, Sheikh IA, Karim S, Haque A, Kamal MA, Chaudhary AG, Azhar E, Mirza Z. 2014 Effect of electromagnetic radiations on neurodegenerative diseases- technological revolution as a curse in disguise. *CNS Neurol Disord Drug Targets* 13:1406-1412.
17. Pall, M. L. 2015 Scientific evidence contradicts findings and assumptions of Canadian Safety Panel 6: microwaves act through voltage-gated calcium channel activation to induce biological impacts at non-thermal levels, supporting a paradigm shift for microwave/lower frequency electromagnetic field action. *Rev. Environ. Health* 3, 99-116. doi: 10.1515/reveh-2015-0001.
18. Bhargav H, Srinivasan TM, Varambally S, Gangadhar BN, Koka P. 2015 Effect of Mobile Phone-Induced Electromagnetic Field on Brain Hemodynamics and Human Stem Cell Functioning: Possible Mechanistic Link to Cancer Risk and Early Diagnostic Value of Electronphotonic Imaging. *J Stem Cells*. 2015;10(4):287-94.
19. Hensinger P, Wilke E. 2016. Mobilfunk-Studienergebnisse bestätigen Risiken Studienrecherche 2016-4 veröffentlicht. *Umwelt Medizin Gesellschaft* 29:3/2016.
20. Houston BJ, Nixon B, King BV, De Iuliis GN, Aitken RJ. 2016 The effects of radiofrequency electromagnetic radiation on sperm function. *Reproduction* 152:R263-R276.
21. Batista Napotnik T, Reberšek M, Vernier PT, Mali B, Miklavčič D. 2016 Effects of high voltage nanosecond electric pulses on eukaryotic cells (in vitro): A systematic review. *Bioelectrochemistry*. 2016 Aug;110:1-12. doi: 10.1016/j.bioelechem.2016.02.011.
22. Asghari A, Khaki AA, Rajabzadeh A, Khaki A. 2016 A review on Electromagnetic fields (EMFs) and the reproductive system. *Electron Physician*. 2016 Jul 25;8(7):2655-2662. doi: 10.19082/2655.
23. Starkey S. 2016 Inaccurate official assessment of radiofrequency safety by the Advisory Group on Non-ionising Radiation. *Rev Environ Health* 31:493-503.
24. Belpomme D, Hardell L, Belyaev I, Burgio E, Carpenter DO. 2018 Thermal and non-thermal health effects of low intensity non-ionizing radiation: An international perspective. *Environ Pollut* 2018 Nov;242(Pt A):643-658. doi: 10.1016/j.envpol.2018.07.019.
25. Pall ML. 2018 How cancer can be caused by microwave frequency electromagnetic field (EMF) exposures: EMF activation of voltage-gated calcium channels (VGCCs) can cause cancer including tumor promotion, tissue invasion and metastasis via 15 mechanisms. Chapter 7 in *Mobile Communications and Public Health*, Marko Markov, Ed., CRC press, pp 167-188.
26. Pall ML. 2018 Wi-Fi is an important threat to human health. *Environ Res* 164:404-416.
27. Wilke I. 2018 Biological and pathological effects of 2.45 GHz on cells, fertility, brain and behavior. *Umwelt Medizin Gesselshaft* 2018 Feb 31 (1).
28. Gautam R, Priyadarshini E, Nirala JP, Rajamani P. 2021. Impact of nonionizing electromagnetic radiation on male infertility: an assessment of the mechanism and consequences. *Int J Radiat Biol* 11,1-22.
29. Lai, H. 2021. Genetic effects of non-ionizing electromagnetic fields. *Electromagn Biol Med*, in press.

**4) Apoptosis/cell death (an important process in production of neurodegenerative diseases that is also important in producing infertility responses) (18 reviews).**

1. Glaser ZR, PhD. 1971 Naval Medical Research Institute Research Report, June 1971. Bibliography of Reported Biological Phenomena (“Effects”) and Clinical Manifestations

- Attributed to Microwave and Radio-Frequency Radiation. Report No. 2 Revised. [https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as\\_sdt=0%2C38](https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as_sdt=0%2C38) (Accessed Sept. 9, 2017)
2. Tolgskaya MS, Gordon ZV. 1973. Pathological Effects of Radio Waves, Translated from Russian by B Haigh. Consultants Bureau, New York/London, 146 pages.
  3. Raines, J. K. 1981. Electromagnetic Field Interactions with the Human Body: Observed Effects and Theories. Greenbelt, Maryland: National Aeronautics and Space Administration 1981; 116 p.
  4. Hardell L, Sage C. 2008. Biological effects from electromagnetic field exposure and public exposure standards. *Biomed. Pharmacother.* 62:104-109. doi: 10.1016/j.biopha.2007.12.004.
  5. Makker K, Varghese A, Desai NR, Mouradi R, Agarwal A. 2009 Cell phones: modern man's nemesis? *Reprod Biomed Online* 18:148-157.
  6. Panagopoulos DJ, Margaritis LH. 2009 Biological and Health Effects of Mobile Telephone Radiations. *Int J Med Biol Front* 2009; 15 (1-2): 33-76.
  7. Levitt, B. B., Lai, H. 2010. Biological effects from exposure to electromagnetic radiation emitted by cell tower base stations and other antenna arrays. *Environ. Rev.* 18, 369-395. doi.org/10.1139/A10-018
  8. Yakymenko I, Sidorik E. 2010 Risks of carcinogenesis from electromagnetic radiation and mobile telephony devices. *Exp Oncol* 32:729-736.
  9. Yakimenko IL, Sidorik EP, Tsybulin AS. 2011 [Metabolic changes in cells under electromagnetic radiation of mobile communication systems]. *Ukr Biokhim Zh* (1999). 2011 Mar-Apr;83(2):20-28.
  10. Pall, ML. 2013. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med* 17:958-965. doi: 10.1111/jcmm.12088.
  11. Kesari KK, Siddiqui MH, Meena R, Verma HN, Kumar S. 2013. Cell phone radiation exposure on brain and associated biological systems. *Ind J Exp Biol* 51: 187-200.
  12. Saliev T, Tachibana K, Bulanin D, Mikhalovsky S, Whitby RD. 2014 Bio-effects of non-ionizing electromagnetic fields in context of cancer therapy. *Front Biosci (Elite Ed)*. 2014 Jan 1;6:175-84. doi: 10.2741/e700.
  13. Pall ML. 2016 Microwave frequency electromagnetic fields (EMFs) produce widespread neuropsychiatric effects including depression. *J Chem Neuroanat* 75(Pt B):43-51. doi: 10.1016/j.jchemneu.2015.08.001.
  14. Batista Napatnik T, Reberšek M, Vernier PT, Mali B, Miklavčič D. 2016 Effects of high voltage nanosecond electric pulses on eukaryotic cells (in vitro): A systematic review. *Bioelectrochemistry*. 2016 Aug;110:1-12. doi: 10.1016/j.bioelechem.2016.02.011.
  15. Asghari A, Khaki AA, Rajabzadeh A, Khaki A. 2016 A review on Electromagnetic fields (EMFs) and the reproductive system. *Electron Physician*. 2016 Jul 25;8(7):2655-2662. doi: 10.19082/2655.
  16. Starkey S. 2016 Inaccurate official assessment of radiofrequency safety by the Advisory Group on Non-ionising Radiation. *Rev Environ Health* 31:493-503.
  17. Pall ML. 2018 Wi-Fi is an important threat to human health. *Environ Res* 164:404-416.
  18. Belyaev I. 2018 Health effects of chronic exposure to radiation from mobile communication. Chapter 4 in: *Mobile Communications and Public Health*, Marko Markov, Ed., CRC Press, pp 65-99.
- 5) **Oxidative stress/free radical damage** (important mechanisms involved in almost all chronic diseases; direct cause of cellular DNA damage). There is published evidence that

oxidative stress is responsible for cumulative changes in tissue structure, producing in turn, much of the cumulative effects of EMF exposure. (29 reviews).

1. Raines, J. K. 1981. *Electromagnetic Field Interactions with the Human Body: Observed Effects and Theories*. Greenbelt, Maryland: National Aeronautics and Space Administration 1981; 116 p.
2. Hardell, L., Sage, C. 2008. Biological effects from electromagnetic field exposure and public exposure standards. *Biomed. Pharmacother.* 62, 104-109.
3. Hazout A, Menezo Y, Madelenat P, Yazbeck C, Selva J, Cohen-Bacrie P. 2008 [Causes and clinical implications of sperm DNA damages]. *Gynecol Obstet Fertil* ;36:1109-1117
4. Makker K, Varghese A, Desai NR, Mouradi R, Agarwal A. 2009 Cell phones: modern man's nemesis? *Reprod Biomed Online* 18:148-157.
5. Desai NR, Kesari KK, Agarwal A. 2009 Pathophysiology of cell phone radiation: oxidative stress and carcinogenesis with focus on the male reproductive system. *Reproduct Biol Endocrinol* 7:114.
6. Yakymenko I, Sidorik E. 2010 Risks of carcinogenesis from electromagnetic radiation and mobile telephony devices. *Exp Oncol* 32:729-736.
7. Kovacic P, Somanathan R. 2010. Electromagnetic fields: mechanisms, cell signaling, other bioprocesses, toxicity, radicals, antioxidants and beneficial effects. *J Recep Sign Transduct* 30:214-226. doi: org/10.3109/10799893.2010.488650
8. Yakymenko IL, Sidorik EP, Tsybulin AS. 2011 [Metabolic changes in cells under electromagnetic radiation of mobile communication systems]. *Ukr Biokhim Zh* (1999). 2011 Mar-Apr;83(2):20-28.
9. Consales, C., Merla, C., Marino, C., et al. 2012. Electromagnetic fields, oxidative stress, and neurodegeneration. *Int. J. Cell Biol.* 2012: 683897.
10. LaVignera et al 2012 La Vignera S, Condorelli RA, Vicari E, D'Agata R, Calogero AE. 2012 Effects of the exposure to mobile phones on male reproduction: a review of the literature. *J Androl* 33:350-356.
11. Pall, ML. 2013. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med* 17:958-965. doi: 10.1111/jcmm.12088.
12. Nazıroğlu M, Yüksel M, Köse SA, Özkaya MO. 2013 Recent reports of Wi-Fi and mobile phone-induced radiation on oxidative stress and reproductive signaling pathways in females and males. *J Membr Biol* 246:869-875.
13. Pall, M. L. 2015. Scientific evidence contradicts findings and assumptions of Canadian Safety Panel 6: microwaves act through voltage-gated calcium channel activation to induce biological impacts at non-thermal levels, supporting a paradigm shift for microwave/lower frequency electromagnetic field action. *Rev. Environ. Health* 3, 99-116.
14. Yakymenko I, Tsybulin O, Sidorik E, Henshel D, Kyrylenko O, Kysylenko S. 2015 Oxidative mechanisms of biological activity of low-intensity radiofrequency radiation. *Electromagnetic Biol Med: Early Online* 1-16. ISSN: 1536-8378.
15. Hensinger P, Wilke E. 2016. Mobilfunk-Studienergebnisse bestätigen Risiken Studienrecherche 2016-4 veröffentlicht. *Umwelt Medizin Gesellschaft* 29:3/2016.
16. Houston BJ, Nixon B, King BV, De Iuliis GN, Aitken RJ. 2016 The effects of radiofrequency electromagnetic radiation on sperm function. *Reproduction* 152:R263-R276.
17. Starkey S. 2016 Inaccurate official assessment of radiofrequency safety by the Advisory Group on Non-ionising Radiation. *Rev Environ Health* 31:493-503.

18. Dasdag S, Akdag MZ. 2016 The link between radiofrequencies emitted from wireless technologies and oxidative stress. *J Chem Neuroanat* 75(Pt B):85-93.
19. Wang H, Zhang X. 2017 Magnetic fields and reactive oxygen species. *Int J Mol Sci*. 2017 Oct 18;18(10). pii: E2175. doi: 10.3390/ijms18102175.
20. Bandara P, Weller S. 2017 Cardiovascular disease: Time to identify emerging environmental risk factors. *Eur J Prevent Cardiol* 24:1819-1823. doi: 10.1177/2047487317734898
21. Pall ML. 2018 Wi-Fi is an important threat to human health. *Environ Res* 164:404-416.
22. Belpomme D, Hardell L, Belyaev I, Burgio E, Carpenter DO. 2018 Thermal and non-thermal health effects of low intensity non-ionizing radiation: An international perspective. *Environ Pollut* 2018 Nov;242(Pt A):643-658. doi: 10.1016/j.envpol.2018.07.019.
23. Belyaev I. 2018 Health effects of chronic exposure to radiation from mobile communication. Chapter 4 in: *Mobile Communications and Public Health*, Marko Markov, Ed., CRC Press, pp 65-99.
24. Yahyazadeh A, Deniz ÖG, Kaplan AA, Altun G, Yurt KK, Davis D. 2018 The genomic effects of cell phone exposure on the reproductive system. *Environ Res*. 2018 Nov;167:684-693. doi: 10.1016/j.envres.2018.05.017.
25. Wilke I. 2018 Biological and pathological effects of 2.45 GHz on cells, fertility, brain and behavior. *Umwelt Medizin Gessellschaft* 2018 Feb 31 (1).
26. Kesari KK, Agarwal A, Henkel R. Radiations and male fertility. 2018 *Reprod Biol Endocrinol*. 2018 Dec 9;16(1):118. doi: 10.1186/s12958-018-0431-1.
27. Santini SJ, Cordone V, Falone S, Mijit M, Tatone C, Amicarelli F, Di Emidio G. 2018 Role of Mitochondria in the Oxidative Stress Induced by Electromagnetic Fields: Focus on Reproductive Systems. *Oxid Med Cell Longev*. 2018 Nov 8;2018:5076271. doi: 10.1155/2018/5076271.
28. Gautam R, Priyadarshini E, Nirala JP, Rajamani P. 2021. Impact of nonionizing electromagnetic radiation on male infertility: an assessment of the mechanism and consequences. *Int J Radiat Biol* 11,1-22.
29. Schuermann D, Mevissen M. 2021. Manmade electromagnetic fields and oxidative stress—biological effects and consequences for health. *Int J Mol Sci* 22: 3772.

**6) Endocrine, that is hormonal effects; Includes changes in non-steroid and also steroid hormones (15 reviews).**

1. Glaser ZR, PhD. 1971 Naval Medical Research Institute Research Report, June 1971. Bibliography of Reported Biological Phenomena (“Effects”) and Clinical Manifestations Attributed to Microwave and Radio-Frequency Radiation. Report No. 2 Revised.  
[https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as\\_sdt=0%2C38](https://scholar.google.com/scholar?q=Glaser+naval+medical+microwave+radio-frequency+1972&btnG=&hl=en&as_sdt=0%2C38) (Accessed Sept. 9, 2017)
2. Tolgskaya MS, Gordon ZV. 1973. *Pathological Effects of Radio Waves*, Translated from Russian by B Haigh. Consultants Bureau, New York/London, 146 pages.
3. Raines, J. K. 1981. *Electromagnetic Field Interactions with the Human Body: Observed Effects and Theories*. Greenbelt, Maryland: National Aeronautics and Space Administration 1981; 116 p.

4. Reiter RJ. 1998 Melatonin in the context of the reported bioeffects of environmental electromagnetic fields. *Bioelectrochemistry & Bioenergetics* 47: 135-142.
  5. Hardell, L., Sage, C. 2008. Biological effects from electromagnetic field exposure and public exposure standards. *Biomed. Pharmacother.* 62, 104-109.
  6. Makker K, Varghese A, Desai NR, Mouradi R, Agarwal A. 2009 Cell phones: modern man's nemesis? *Reprod Biomed Online* 18:148-157.
  7. Gye MC, Park CJ. 2012 Effect of electromagnetic field exposure on the reproductive system. *Clin Exp Reprod Med* 39:1-9. doi.org/10.5653/cerm.2012.39.1.1
  8. Halgamuge MN. 2013 Pineal melatonin level disruption in humans due to electromagnetic fields and ICNIRP limits. *Radiat Prot Dosimetr* 154: 405-416
  9. Pall, M. L. 2015. Scientific evidence contradicts findings and assumptions of Canadian Safety Panel 6: microwaves act through voltage-gated calcium channel activation to induce biological impacts at non-thermal levels, supporting a paradigm shift for microwave/lower frequency electromagnetic field action. *Rev. Environ. Health* 3, 99-116.
  10. Sangün Ö, Dündar B, Çömlekçi S, Büyükgebiz A. 2016 The Effects of Electromagnetic Field on the Endocrine System in Children and Adolescents. *Pediatr Endocrinol Rev* 13:531-545.
  11. Hecht, Karl. 2016 Health Implications of Long-Term Exposures to Electrosmog. Brochure 6 of A Brochure Series of the Competence Initiative for the Protection of Humanity, the Environment and Democracy. [http://kompetenzinitiative.net/KIT/wp-content/uploads/2016/07/KI\\_Brochure-6\\_K\\_Hecht\\_web.pdf](http://kompetenzinitiative.net/KIT/wp-content/uploads/2016/07/KI_Brochure-6_K_Hecht_web.pdf) (accessed Feb. 11, 2018)
  12. Asghari A, Khaki AA, Rajabzadeh A, Khaki A. 2016 A review on Electromagnetic fields (EMFs) and the reproductive system. *Electron Physician*. 2016 Jul 25;8(7):2655-2662. doi: 10.19082/2655.
  13. Warille AA, Altun G, Elamin AA, Kaplan AA, Mohamed H, Yurt KK, Elhaj AE. 2017 Skeptical approaches concerning exposure to electromagnetic fields on brain hormones and enzyme activities. *J Microscopy Ultrastruct* 5:177-184. <https://doi.org/10.1016/j.jmau.2017.09.002>
  14. Pall ML. 2018 Wi-Fi is an important threat to human health. *Environ Res* 164:404-416.
  15. Wilke I. 2018 Biological and pathological effects of 2.45 GHz on cells, fertility, brain and behavior. *Umwelt Medizin Gessellschaft* 2018 Feb 31 (1).
- 7) **Increased intracellular calcium levels, thought to be the cause in all other effects** (16 different reviews).
1. Adey WR. 1988 Cell membranes: the electromagnetic environment and cancer promotion. *Neurochem Res*.13:671-677.
  2. Walleczek, J. 1992. Electromagnetic field effects on cells of the immune system: the role of calcium signaling. *FASEB J.* 6, 3177-3185.
  3. Adey, WR. 1993 Biological effects of electromagnetic fields. *J Cell Biochem* 51:410-416.
  4. Frey AH. 1993 Electromagnetic field interactions with biological systems. *FASEB J* 7:272-281.
  5. Funk RHW, Monsees T, Özkucur N. 2009 Electromagnetic effects—From cell biology to medicine. *Prog Histochem Cytochem* 43:177-264.

6. Yakymenko IL, Sidorik EP, Tsybulin AS. 1999 [Metabolic changes in cells under electromagnetic radiation of mobile communication systems]. Ukr Biokhim Zh (1999), 2011 Mar-Apr:20-28.
  7. Gye MC, Park CJ. 2012 Effect of electromagnetic field exposure on the reproductive system. Clin Exp Reprod Med 39:1-9. doi.org/10.5653/cerm.2012.39.1.1
  8. Pall, ML. 2013. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. J Cell Mol Med 17:958-965. doi: 10.1111/jcmm.12088.
  9. Pall ML. 2014 Electromagnetic field activation of voltage-gated calcium channels: role in therapeutic effects. Electromagn Biol Med. 2014 Apr 8 doi: 10.3109/15368378.2014.906447.
  10. Pall ML. 2015 How to approach the challenge of minimizing non-thermal health effects of microwave radiation from electrical devices. International Journal of Innovative Research in Engineering & Management (IJIREM) ISSN: 2350-0557, Volume-2, Issue -5, September 2015; 71-76.
  11. Pall, M. L. 2015 Scientific evidence contradicts findings and assumptions of Canadian Safety Panel 6: microwaves act through voltage-gated calcium channel activation to induce biological impacts at non-thermal levels, supporting a paradigm shift for microwave/lower frequency electromagnetic field action. Rev. Environ. Health 3, 99-116. doi: 10.1515/reveh-2015-0001.
  12. Pall ML. 2016 Electromagnetic fields act similarly in plants as in animals: Probable activation of calcium channels via their voltage sensor. Curr Chem Biol 10: 74-82.
  13. Pall ML. 2016 Microwave frequency electromagnetic fields (EMFs) produce widespread neuropsychiatric effects including depression. J Chem Neuroanat 75(Pt B):43-51. doi: 10.1016/j.jchemneu.2015.08.001.
  14. Batista Napotnik T, Reberšek M, Vernier PT, Mali B, Miklavčič D. 2016 Effects of high voltage nanosecond electric pulses on eukaryotic cells (in vitro): A systematic review. Bioelectrochemistry. 2016 Aug;110:1-12. doi: 10.1016/j.bioelechem.2016.02.011.
  15. Asghari A, Khaki AA, Rajabzadeh A, Khaki A. 2016 A review on electromagnetic fields (EMFs) and the reproductive system. Electron Physician. 2016 Jul 25;8(7):2655-2662. doi: 10.19082/2655.
  16. Kesari KK, Agarwal A, Henkel R. Radiations and male fertility. 2018 Reprod Biol Endocrinol. 2018 Dec 9;16(1):118. doi: 10.1186/s12958-018-0431-1.
- 8) **Cancer including initiation, promotion and progression, further including tumor progression, tissue invasion and metastasis** (44 reviews). I am going to attach a copy of my paper, #37 below which shows how the primary effect of EMF exposures, activation of voltage-gated calcium channels (VGCCs) can act to elevate 15 different mechanisms known to have roles in carcinogenesis, the process of cancer causation. That paper shows that 15 mechanisms produced following EMF exposure, each have roles in causing cancer.
1. Dwyer, M. J., Leeper, D. B. 1978 A Current Literature Report on the Carcinogenic Properties of Ionizing and Nonionizing Radiation. DHEW Publication (NIOSH) 78-134, March 1978.
  2. Marino AA, Morris DH. 1985 Chronic electromagnetic stressors in the environment. A risk factor in human cancer. J environ sci health C3:189-219.

3. Adey WR. 1988 Cell membranes: the electromagnetic environment and cancer promotion. *Neurochem Res.*13:671-677.
4. Adey WR. 1990 Joint actions of environmental nonionizing electromagnetic fields and chemical pollution in cancer promotion. *Environ Health Perspect* 86:297-305.
5. Frey AH. 1993 Electromagnetic field interactions with biological systems. *FASEB J* 7:272-281.
6. Goldsmith JR. 1995 Epidemiological evidence of radiofrequency radiation (microwave) effects on health in military, broadcasting and occupational settings. *Int J Occup Environ Health* 1:47-57.
7. Goldsmith JR. 1997 Epidemiologic evidence relevant to radar (microwave) effects. *Env Health Perspect* 105(Suppl 6):1579-1587.
8. Goldsmith JR. 1997 TV broadcast towers and cancer: the end of innocence for radiofrequency exposures. *Am J Ind Med* 32:689-692.
10. Stepansky R, Jahn O, Windischbauer G, Zeitlhofer J. 2000 [Electromagnetic fields--effects on health]. *Acta Med Austriaca.* 2000;27(3):69-77.
9. Kundi M, Mild K, Hardell L, Mattsson M. 2004 Mobile telephones and cancer – a review of the epidemiological evidence. *J Toxicol Env Health, Part B* 7:351-384.
10. Kundi M. 2004 Mobile phone use and cancer. *Occup Env Med* 61:560-570.
- Oberfeld, G., Navarro, A.E., Portoles, M., Maestu, C., Gomez-Perretta, C., 2004. The microwave syndrome: further aspects of a Spanish study, <http://www.apdr.info/electrocontaminacion/Documentos/Investigacion/ESTUDOS%20EPIDEMIOLOXIDOS%20E%20ANTENAS/The%20Microwave%20Syndrome%20-%20Further%20Aspects%20of%20a%20Spanish%20Study.pdf>.
11. Behari J, Paulraj R. 2007 Biomarkers of induced electromagnetic field and cancer. *Indian J Exp Biol* 45:77-85.
12. Hardell L, Carlberg M, Soderqvist F, Hansson Mild K. 2008 Meta-analysis of long-term mobile phone use and the association with brain tumors. *Int J Oncol* 32:1097-1103.
13. Khurana VG, Teo C, Kundi M, Hardell L, Carlberg M. 2009 Cell phones and brain tumors: a review including the long-term epidemiologic data. *Surg Neurol* 72:205-214.
14. Desai NR, Kesari KK, Agarwal A. 2009 Pathophysiology of cell phone radiation: oxidative stress and carcinogenesis with focus on the male reproductive system. *Reproduct Biol Endocrinol* 7:114.
15. Davanipour Z, Sobel E. 2009 Long-term exposure to magnetic fields and the risks of Alzheimer's disease and breast cancer: Further biological research. *Pathophysiology* 16:149-156.
16. Yakymenko I, Sidorik E. 2010 Risks of carcinogenesis from electromagnetic radiation and mobile telephony devices. *Exp Oncol* 32:729-736.
17. Carpenter DO. 2010 Electromagnetic fields and cancer: the cost of doing nothing. *Rev Environ Health* 25:75-80.
18. Giuliani L, Soffriti M (Eds). 2010 NON-THERMAL EFFECTS AND MECHANISMS OF INTERACTION BETWEEN ELECTROMAGNETIC FIELDS AND LIVING MATTER, RAMAZZINI INSTITUTE EUR. J. ONCOL. LIBRARY Volume 5, National Institute for the Study and Control of Cancer and Environmental Diseases “Bernardino Ramazzini” Bologna, Italy 2010, 400 page monograph.
19. Khurana, V. G., Hardell, L., Everaert, J., Bortkiewicz, A., Carlberg, M., Ahonen, M. 2010 Epidemiological evidence for a health risk from mobile phone base stations. *Int. J. Occup. Environ. Health* 16, 263-267.

20. Yakymenko, I., Sidorik, E., Kyrylenko, S., Chekhun, V. 2011. Long-term exposure to microwave radiation provokes cancer growth: evidences from radars and mobile communication systems. *Exp. Oncol.* 33(2), 62-70.
21. Bioinitiative Working Group, David Carpenter and Cindy Sage (eds). 2012 Bioinitiative 2012: A rationale for biologically-based exposure standards for electromagnetic radiation. <http://www.bioinitiative.org/participants/why-we-care/>
22. Ledoigt G, Belpomme D. 2013 Cancer induction molecular pathways and HF-EMF irradiation. *Adv Biol Chem* 3:177-186.
23. Hardell L, Carlberg M. 2013 Using the Hill viewpoints from 1965 for evaluating strengths of evidence of the risk for brain tumors associated with use of mobile and cordless phones. *Rev Environ Health* 28:97-106. doi: 10.1515/reveh-2013-0006.
24. Hardell L, Carlberg M, Hansson Mild K. 2013 Use of mobile phones and cordless phones is associated with increased risk for glioma and acoustic neuroma. *Pathophysiology* 2013;20(2):85-110.
25. Carpenter DO. 2013 Human disease resulting from exposure to electromagnetic fields. *Rev Environ Health* 2013;28:159-172.gj
26. Davis DL, Kesari S, Soskolne CL, Miller AB, Stein Y. 2013 Swedish review strengthens grounds for concluding that radiation from cellular and cordless phones is a probable human carcinogen. *Pathophysiology* 20:123-129.
27. Morgan LL, Miller AB, Sasco A, Davis DL. 2015 Mobile phone radiation causes brain tumors and should be classified as a probable human carcinogen (2A). *Int J Oncol* 46(5): 1865-1871.
28. Mahdavi M, Yekta R, Tackallou SH. 2015 Positive correlation between ELF and RF electromagnetic fields on cancer risk. *J Paramed Sci* 6(3), ISSN 2008-4978.
29. Carlberg M, Hardell L. 2017 Evaluation of Mobile Phone and Cordless Phone Use and Glioma Risk Using the Bradford Hill Viewpoints from 1965 on Association or Causation. *BioMed Res Int* 2017, Article ID 9218486, <https://doi.org/10.1155/2017/9218486>
30. Bortkiewicz A, Gadzicka E, Szymczak W. 2017 Mobile phone use and risk for intracranial tumors and salivary gland tumors - A meta-analysis. *Int J Occup Med Environ Health* 30:27-43.
31. Bielsa-Fernández P, Rodríguez-Martín B. 2017 [Association between radiation from mobile phones and tumour risk in adults]. *Gac Sanit.* 2017 Apr 12. pii: S0213-9111(17)30083-3. doi: 10.1016/j.gaceta.2016.10.014. [Epub ahead of print]
32. Alegría-Loyola MA, Galnares-Olalde JA, Mercado M. 2017 [Tumors of the central nervous system]. *Rev Med Inst Mex Seguro Soc* 55:330-334.
33. Yang M, Guo W, Yang C, Tang J, Huang Q, Feng S, Jiang A, Xu X, Jiang G. 2017. Mobile phone use and glioma risk: A systematic review and meta-analysis. *PLoS One.* 2017 May 4;12(5):e0175136.
34. Prasad M, Kathuria P, Nair P, Kumar A, Prasad K. 2017 Mobile phone use and risk of brain tumours: a systematic review of association between study quality, source of funding, and research outcomes. *Neurol Sci.* 2017 Feb 17. doi: 10.1007/s10072-017-2850-8. [Epub ahead of print].
35. Miller A. 2017 References on cell phone radiation and cancer. <https://ehtrust.org/references-cell-phone-radio-frequency-radiation-cancer/> (Accessed Sept. 9, 2017)
36. Hardell L. 2017 World Health Organization, radiofrequency radiation and health – a hard nut to crack (Review). *Int J Oncol* 51:405-413.
37. Belpomme D, Hardell L, Belyaev I, Burgio E, Carpenter DO. 2018 Thermal and non-thermal health effects of low intensity non-ionizing radiation: An international



perspective. *Environ Pollut* 2018 Nov;242(Pt A):643-658. doi: 10.1016/j.envpol.2018.07.019.

38. Pall ML. 2018 How cancer can be caused by microwave frequency electromagnetic field (EMF) exposures: EMF activation of voltage-gated calcium channels (VGCCs) can cause cancer including tumor promotion, tissue invasion and metastasis via 15 mechanisms. Chapter 7 in: *Mobile Communications and Public Health*, Marko Markov, Ed., CRC Press, pp 167-188.

39. Subhan F, Khan A, Ahmed S, Malik SN, Bakshah ST, Tahir S.. 2018 Mobile antenna's and its impact on human health. *J Medical Imaging and Health Informatics*. 8(66):11266 - 1273. doi: 10.1166/jjmihi.22018.2296.

40. Belyaev I. 2018 Health effects of chronic exposure to radiation from mobile communication. Chapter 4 in: *Mobile Communications and Public Health*, Marko Markov, Ed., CRC Press, pp 65-99.

41. Kocaman A, Altun G, Kaplan AA, Deniz OG, Yurt KK, Kaplan S. 2018 Genotoxic and carcinogenic effects of non-ionizing electromagnetic fields. *Environ Res* 163: 71-79. doi: 10.1016/j.envres.2018.01.034

42. Miller AB, Morgan LL, Udasin I, Davis DL. 2018 Cancer epidemiology update, following the 2011 IARC evaluation of radiofrequency electromagnetic fields (Monograph 102). *Environmental Research*. 167:673-683. Nov 2018. doi: 10.1016/j.envres.2018.06.043.

43. Mialon HM., Nesson ET. 2020. The association between mobile phones and the risk of brain cancer mortality: A 25-year cross-country analysis. *Contemp Econ Policy* 38:258-269.

44. Choi Y-J, Moskowitz JM, Myung S-K, Lee Y-R, Hong Y-C. 2020 Cellular Phone Use and Risk of Tumors: Systematic Review and Meta-Analysis. *International Journal of Environmental Research and Public Health*. 2020, 17(21), 8079.

**(9). There are also 15 additional reviews on cardiac effects. Those cardiac effects, include tachycardia, arrhythmia and bradycardia (with bradycardia typically reported after long times of exposures). Some recent studies have also reported heart palpitations. Arrhythmias, especially when they are associated with either bradycardia or severe tachycardia, are often associated with sudden cardiac death. Sudden cardiac death causes over 5% of the total mortality in technologically advanced countries, so this could be a major source of EMF-caused fatality.**

1. Raines, J. K. 1981. *Electromagnetic Field Interactions with the Human Body: Observed Effects and Theories*. Greenbelt, Maryland: National Aeronautics and Space Administration 1981; 116 p.
2. Frey AH, Eichert ES. 1986 Modification of heart function with low intensity electromagnetic energy. *J Bioelect* 5:201-210. (Note: this is a primary literature citation that also reviews some important earlier studies)
3. Jauchem JR. 1997. Exposure to extremely-low-frequency electromagnetic fields and radiofrequency radiation: cardiovascular effects in humans. *Int Arch Occup Environ Health* 70: 9-21.
4. Pakhomov AG, Akyel Y, Pakhomova ON, Stuck BE, Murphy MR. Current state and implications of research on biological effects of millimeter waves: a review of the literature. *Bioelectromagnetics* 1998;19:393-413.

5. Cherry N. 2002 Cardiac Effects of Natural and Artificial EMR. [http://www.neilcherry.nz/documents/90\\_s4\\_EMR\\_Cardiac\\_Effects\\_Review.pdf](http://www.neilcherry.nz/documents/90_s4_EMR_Cardiac_Effects_Review.pdf) (Accessed August 3, 2019).
6. Scherling BJ, Yamanashi WS, Hou Y, Jacobson JI, Jackman WM, Lazzara R. Magnetism and cardiac arrhythmias. *Cardiol Rev* 2004;12:85-96.
7. Hardell, L., Sage, C. 2008. Biological effects from electromagnetic field exposure and public exposure standards. *Biomed. Pharmacother.* 62, 104-109.
8. Consensus paper of the Austrian Medical Association's EMF working group. 2012 Guideline of the Austrian Medical Association for the diagnosis and treatment of EMF-related health problems and illnesses (EMF syndrome). <https://www.vagbrytaren.org/Guideline%20%20AG-EMF.pdf> (Accessed 01/2019).
9. Havas M. 2013 Radiation from wireless technology affects the blood, the heart and the autonomic nervous system. *Rev Environ Health* 28: 75-84.
10. Sage C. 2015 The implications of non-linear biological oscillations on human electrophysiology for electrohypersensitivity (EHS) and multiple chemical sensitivity (MCS). *Rev Environ Health*. 2015;30(4):293-303. doi: 10.1515/reveh-2015-0007.
11. Hecht, Karl. 2016 Health Implications of Long-Term Exposures to Electrosmog. Brochure 6 of A Brochure Series of the Competence Initiative for the Protection of Humanity, the Environment and Democracy. [http://kompetenzinitiative.net/KIT/wp-content/uploads/2016/07/KI\\_Brochure-6\\_K\\_Hecht\\_web.pdf](http://kompetenzinitiative.net/KIT/wp-content/uploads/2016/07/KI_Brochure-6_K_Hecht_web.pdf) (accessed Feb. 11, 2018)
12. Belyaev I, Dean A, Eger H, Hubmann G, Jandrisovits R, Kern M, Kundi M, Moshammer H, Lercher P, Müller K, Oberfeld G, Ohnsorge P, Pelzmann P, Scheingraber C, Thill R. 2016 EUROPAEM EMF Guideline 2016 for the prevention, diagnosis and treatment of EMF-related health problems and illnesses. *Rev Environ Health* DOI 10.1515/reveh-2016-0011.
13. Wilke I. 2018 Biological and pathological effects of 2.45 GHz on cells, fertility, brain and behavior. *Umwelt Medizin Gesellschaft* 2018 Feb 31 (1).
14. Subhan F, Khan A, Ahmed S, Malik SN, Bakshah ST, Tahir S. 2018 Mobile antenna's and its impact on human health. *J Medical Imaging and Health Informatics*. 8(66):11266 - 1273. doi: 10.1166/jjmihi.22018.2296.
15. Pall ML. 2021. Millimeter (MM) wave and microwave frequency radiation produce deeply penetrating effects: the biology and the physics. *Rev Environ Health*, vol. , no. , 2021, pp. 000010151520200165. <https://doi.org/10.1515/reveh-2020-0165>

We have here, a total of 214 bodies of evidence each showing that non-thermal exposures well below ICNIRP, FCC or other “safety guidelines” cause an important health-related effect. These 9 different non-thermal effects are not the only effects apparently being produced. These 214 bodies of evidence *individually* provide strong and in many cases compelling evidence against any claims that can be made on the basis of the ICNIRP, US FCC, EU or other similar “safety guidelines” as do thousands of primary literature citations. There are fewer than 214 individual reviews listed above because many of these reviews provide bodies of evidence for multiple EMF effects.

What response do we have from the telecommunications industry or other organizations that have supported the industry positions. The only thing that we have are statements similar to the statement put out by Dr. Jeffrey Shuren, M.D., J.D., Director of the FDA's Center for Devices and Radiological Health on the National Toxicology Program's report on radiofrequency energy exposure as follows: "Based on our ongoing evaluation of this issue, the *totality of the available scientific evidence* (italics added) continues to not support adverse health effects in humans caused by exposures at or under the current radiofrequency energy exposure limits. We believe the existing safety limits for cell phones remain acceptable for protecting the public health." This

statement, as were similar statements from other organizations, provided not one iota of evidence in support of the claims. This is obviously unacceptable. What such individuals and organizations need to do when considering the *totality of the evidence* is to consider each of the reviews providing evidence for each of these nine different effects, as well as each of the relevant underlying primary literature citations cited in these reviews. If they wish to rebut these repeated findings, they need to cite each of these reviews, providing an objective description of the relevant evidence described within them and then and only then, provide whatever evidence they may have rebutting the positions taken in these reviews. The complete failure to do this means that the positions taken by Dr. Shuren and similar positions of others on this are fatally flawed. Those flaws go to the heart of the scientific method. Dr. Karl Popper, one of the two most important philosophers of science of the 20<sup>th</sup> century, has argued compellingly that falsifying information, information that falsifies a theory or hypothesis, is the most important type of information in science. Here we have 213 bodies of evidence that falsify the position of the industry and organizations that historically have supported the industry with no response whatsoever other than a completely undocumented denial.

2. There were also 13 reviews cited in Chapter 1 of my 90 page document and listed immediately below, 10 of which were also cited in my 2018 Wi-Fi paper, each of which showed that pulsed EMFs are, in most cases, much more biologically active than are non-pulsed (also known as continuous wave) EMFs of the same average intensity. The pulses here, are modulating pulses, modulating the intensity of an underlying frequency. Because average intensities, averaged over a 6 minute period or even worse a 30 minute period, are the basis of the ICNIRP, US FCC, EU, SCENIHR and Canadian guidelines this is a fatal flaw in the structure of those safety guidelines. Average intensities are *not* predictive of biological effects and therefore cannot be used as the basis of any useful regulatory scheme. Pulsation is also of great importance, because all wireless communication devices, with the exception of FM radio communication, communicate at least in part, via modulating pulsation and the smarter they are, the more they pulse. Radar units also expose us to pulsations because of the phased arrays that are involved. Consequently, the role of pulsation is stunningly important with regard to the EMFs we are most exposed to.

13 Reviews Each Showing that Pulsed EMFs Are, in Most Cases Much More Biologically Active than Are Non-Pulsed (Continuous Wave) EMFs of the Same Average Intensity:

1. Osipov YuA, 1965 [Labor hygiene and the effect of radiofrequency electromagnetic fields on workers]. Leningrad Meditsina Publishing House, 220 pp.
2. Pollack H, Healer J. 1967 Review of Information on Hazards to Personnel from High-Frequency Electromagnetic Radiation. Institute for Defense Analyses; Research and Engineering Support Division. IDA/HQ 67-6211, Series B, May 1967.
3. Frey AH. 1974 Differential biologic effects of pulsed and continuous electromagnetic fields and mechanisms of effect. Ann N Y Acad Sci 238: 273-279.
4. Creighton MO, Larsen LE, Stewart-DeHaan PJ, Jacobi JH, Sanwal M, Baskerville JC, Bassen HE, Brown DO, Trevithick JR. 1987 In vitro studies of microwave-induced cataract. II. Comparison of damage observed for continuous wave and pulsed microwaves. Exp Eye Res 45:357-373.
5. Grigor'ev IuG. 1996 [Role of modulation in biological effects of electromagnetic radiation]. Radiats Biol Radioecol 36:659-670.
6. Belyaev I. 2005 Non-thermal biological effects of microwaves. Microwave Rev 11:13-29.
7. Belyaev I. 2005 Non-thermal biological effects of microwaves: current knowledge, further perspective and urgent needs. Electromagn Biol Med 24(3):375-403.

8. Markov MS. 2007 Pulsed electromagnetic field therapy: History, state of the art and future. *The Environmentalist* 27:465-475.
9. Van Boxem K, Huntoon M, Van Zundert J, Patijn J, van Kleef M, Joosten EA. 2014 Pulsed radiofrequency: a review of the basic science as applied to the pathophysiology of radicular pain: a call for clinical translation. *Reg Anesth Pain Med.* 2014 Mar-Apr;39(2):149-59.
10. Belyaev, I. 2015. Biophysical mechanisms for nonthermal microwave effects. In: *Electromagnetic Fields in Biology and Medicine*, Marko S. Markov, ed, CRC Press, New York, pp 49-67.
11. Pall, M. L. 2015 Scientific evidence contradicts findings and assumptions of Canadian Safety Panel 6: microwaves act through voltage-gated calcium channel activation to induce biological impacts at non-thermal levels, supporting a paradigm shift for microwave/lower frequency electromagnetic field action. *Rev. Environ. Health* 3, 99-116. doi: 10.1515/reveh-2015-0001.
12. Panagopoulos DJ, Johansson O, Carlo GL. 2015 Real versus simulated mobile phone exposures in experimental studies. *BioMed. Res. Int.* 2015, article ID 607053, 8 pages. doi: 10.1155/2015/607053.
13. Batista Napotnik T, Reberšek M, Vernier PT, Mali B, Miklavčič D. 2016 Effects of high voltage nanosecond electric pulses on eukaryotic cells (in vitro): A systematic review. *Bioelectrochemistry.* 2016 Aug;110:1-12. doi: 10.1016/j.bioelechem.2016.02.011.

3. The following comes from (Pall ML. 2018 Wi-Fi is an important threat to human health. *Environ Res.* 2018 Jul;164:405-416; full citations can be obtained from that paper; information inserted into that text is italicized to identify it): How are the non-thermal EMF effects produced? The author found the answer to this question in the already published scientific literature (Pall, 2013). That study showed that in 24 different studies [there are now a total of 26 Pall, (2015b) *and two additional examples were cited in the Wi-Fi paper, for a total of 28*], effects of low-intensity EMFs, including microwave frequency and also extremely low frequency EMFs, static electrical fields and static magnetic fields could be blocked by calcium channel blockers, drugs that are specific for blocking voltage-gated calcium channels (VGCCs). There were 5 different types of calcium channel blockers used in these studies, each thought to be highly specific, each structurally distinct and each binding to a different site on the VGCCs. In papers where multiple effects were studied, all studied effects were blocked or greatly lowered by calcium channel blockers. These studies show that EMFs produce diverse non-thermal effects via VGCC activation Pall 2013; 2014; 2015a & b; 2016a & b) in many human and animal cells. In plant cells, EMFs activate somewhat similar calcium channels and produce somewhat similar effects on oxidative stress, cellular DNA damage and calcium signaling (Pall, 2016a). Furthermore, many different effects shown to be produced in repeated studies by EMF exposures, including the effects discussed above, can be produced by downstream effects of VGCC activation, via increased [Ca<sup>2+</sup>]<sub>i</sub>, as discussed in detail below.

Before leaving this issue, it is important to discuss why the VGCCs are so sensitive to activation by these low-intensity EMFs. The VGCCs each have a voltage sensor which is made up of 4 alpha helices in the plasma membrane, with each such helix having 5 positive charges on it, for a total of 20 positive charges (Pall, 2015b). These voltage sensor helices are each called S4 helices because each is the fourth helix in a distinct multi-helix domain. Each of these voltage sensor charges is within the lipid bilayer part of the plasma membrane. The electrical forces on the voltage sensor are very high for three distinct reasons (Pall 2015b; 2015a; 2016a). 1. The 20 charges on the voltage sensor make the forces on voltage sensor 20 times higher than the forces on a single charge. 2. Because these charges are within the lipid bilayer section of the membrane where the dielectric constant is about 1/120th of the dielectric constant of the aqueous parts of the

cell, the law of physics called Coulomb's law, predicts that the forces on those charges will be approximately 120 times higher than the forces on charges in the aqueous parts of the cell. 3. Because the plasma membrane has a high electrical resistance whereas the aqueous parts of the cell are highly conductive, the electrical gradient across the plasma membrane is estimated to be concentrated about 3000-fold, as shown by Ohm's law. The combination of these effects means that comparing the forces on the voltage sensor with the forces on singly charged groups in the aqueous parts of the cell, the forces on the voltage sensor are approximately  $20 \times 120 \times 3000 = 7.2$  million times higher (Pall, 2015b). The physics predicts, therefore, extraordinarily strong forces activating the VGCCs via the voltage sensor. It follows that the biology tells us that the VGCCs are the main target of the EMFs and the physics tells us why they are the main target. Thus the physics and biology are pointing in the same direction. All of these findings contradict the basic assumptions of the safety guidelines which are based on average exposures averaged over at least 6 minutes and which set allowable levels based on SAR, a measure of tissue heating and therefore are only relevant to thermal effects.

There are also additional findings pointing to the voltage sensor as the direct target of the EMFs. In addition to the VGCCs, there are also voltage-gated sodium, potassium and chloride channels, with each of these having a voltage sensor similar to those found in the VGCCs. Lu et al (2015) reported that voltage gated sodium channels, in addition to the VGCCs were activated by EMFs. Tabor et al (2014) found that Mauthner cells, specialized neurons with special roles in triggering rapid escape mechanisms in fish, were almost instantaneously activated by electrical pulses, which acted via voltage-gated sodium channel activation to subsequently produce large  $[Ca^{2+}]_i$  increases. Zhang et al (2016) reported that in addition to the VGCCs, potassium and chloride channels were each activated by EMFs, although these other voltage-gated ion channels had relatively modest roles compared with the VGCCs in producing biological effects. Each of these three studies, the Lu et al (2015) study, the Tabor et al (2014) study and the Zhang et al (2016) study used specific blockers for these other voltage-gated ion channels to determine their roles. The Tabor et al (2014) study also used genetic probing to determine the role of the voltage-gated sodium channels. Lu et al (2015) also used whole cell patch clamp measurements to measure the rapid influx of both sodium and calcium into the cell via the voltage-gated channels following EMF exposure. *One important finding that is not in the Wi-Fi paper is that Tekieh et al, in a 2016 paper (Effects of electromagnetic field exposure on conduction and concentration of voltage gated calcium channels: A Brownian dynamics study. Brain Res. 2016 Sep 1;1646:560-569), showed that VGCCs in isolated plasma membranes, were activated by three different frequencies of microwave radiation. That shows that EMF activation of the VGCCs is directly produced by EMF impact on the VGCC protein.* Sodium influx, particularly in electrically active cells, act in the normal physiology to depolarize the plasma membrane, leading to VGCC activation such that the voltage-gated sodium channels may act primarily via indirect activation of the VGCCs. In summary then, we have evidence that in animal including human cells, seven distinct classes of voltage-gated ion channels are each activated by EMF exposures: From the Pall, 2013 review, four classes of voltage-gated ion channels were shown from calcium channel blocker studies, to be activated by EMFs, L-type, T-type, N-type and P/Q -type VGCCs. In this paragraph we have evidence that three other channels are also activated, voltage-gated sodium channels, voltage-gated potassium channels and voltage-gated chloride channels. Furthermore the plant studies strongly suggest that the so called TPC channels, which contain a similar voltage sensor, are activated in plants allowing calcium influx into plants to produce similar EMF-induced responses (Pall 2016a). One can put those observations together with the powerful findings from the physics, that the electrical forces on the voltage-sensor are stunningly strong, something like 7.2 million times stronger than the forces on the singly charged groups in the aqueous phases of the cell. Now you have a stunningly powerful argument that the voltage sensor is the predominant direct target of the EMFs. *Because heating is produced predominantly by the EMF forces on*

*singly electrically charged groups in aqueous solution, the 7.2 million figure suggests that safety guidelines allow us to be exposed to EMFs that are approximately 7.2 million times too high. The failure of the "safety guidelines" to discuss the relevant physics of the voltage sensor means that the physics underlying the "safety guidelines" is deeply flawed.*

There is one additional finding that should be discussed here. In a study published by Pilla (2012), it was found that pulsed EMFs produced an "instantaneous" increase in calcium/calmodulin-dependent nitric oxide synthesis in cells in culture. What Pilla (2012) showed was that following EMF exposure, the cells in culture, must have produced a large increase in  $[Ca^{2+}]_i$ , this in turn produced a large increase in nitric oxide synthesis, the nitric oxide diffused out of the cells and out of the aqueous medium above the cells into the gas phase, where the nitric oxide was detected by a nitric oxide electrode. This entire sequence occurred in less than 5 seconds. This eliminates almost any conceivable indirect effect, except possibly via plasma membrane depolarization. Therefore that the pulsed EMFs are acting directly on the voltage sensors of the VGCCs and possibly the voltage-gated sodium channels, to produce the  $[Ca^{2+}]_i$  increase.

Why is it that the VGCCs, acting via calcium influx, seem to be much more important in producing EMF effects than are the other voltage-gated ion channels? Probably for three reasons: 1.  $Ca^{2+}$  ions under resting conditions in cells have about a 10,000-fold concentration gradient driving them into the cell, and over a million-fold electrochemical gradient also driving them into the cell. Because of this, one can have huge calcium influxes upon channel activation. 2.  $[Ca^{2+}]_i$  produces many important regulatory effects, such that over activation of those effects can have very large pathophysiological consequences. 3. Sustained elevation of  $[Ca^{2+}]_i$  produces major cell damage.

*This section of the Wi-Fi paper was followed by an additional section showing how VGCC activation acting via elevated  $[Ca^{2+}]_i$ , can produce each of the non-thermal effects documented above and elsewhere in the scientific literature.*

**4 & 5.** There is a large literature on nanosecond pulses producing biological effects. If you search under nanosecond pulse in the EMF-Portal database, you will find 213 hits where when each of these are examined individually, over 100 are genuine nanosecond pulse studies that produced non-thermal effects. These do produce effects but when they these pulses have their intensities averaged over 6 minutes or 30 minutes, that fall far short of the levels that "safety guidelines" predict are needed to produce effects. This discrepancy with "safety guidelines" was noted in the second earliest nanosecond pulse study listed in the EMF-portal database [Raslear TG, Akyel Y, Bates F, Belt M, Lu ST. 1993. Temporal bisection in rats: the effects of high-peak-power pulsed microwave irradiation. *Bioelectromagnetics* 14:459-478]. If you take a typical pulse that may last for let say 40 nanoseconds and average it over a 6 minute period (about  $10^{10}$  times longer), as the FCC, EU and other "safety guidelines" do, the average intensity (and average SAR) is so low that, of course, the safety guidelines predict there cannot be effects. ***But there are repeatedly found effects in nanosecond pulses ranging from 2 ns to 600 ns.*** So here again the "safety guidelines" are not predictive of biological effects. It makes no sense to average intensities over approximately  $10^{10}$  times longer than it takes to produce an effect. The logic here is the same as if the following were to occur: Let's assume that you are concerned about someone shooting you with a high power rifle bullet traveling at about 700 meters per second. The bullet takes about 50 microseconds to tear your body apart. If someone from a regulatory authority tells you that you don't need to worry about that, if you average the force of the rifle bullet over a 21 day period (about  $10^{10}$  times longer than 50 microseconds), the average intensity is so low, you don't need to worry about it. If someone were to tell you that, you would laugh in their face and state that they

are either completely incompetent or completely corrupt. That is exactly the correct response in dealing with the EMF safety guidelines of the regulatory authorities. There are several of these nanosecond pulse studies that have shown that VGCC activation has a key role in producing them [Azarov et al, 2019 Excitation of murine cardiac myocytes by nanosecond pulsed electric field. *J Cardiovasc Electrophys* 30:392-401; Hristov et al, 2018. Expression of voltage-gated calcium channels augments cell susceptibility to membrane disruption by nanosecond pulsed electric fields. *Biochim Biophys Acta Biomembr* 1860:2175-2183; Vernier PT, Sun Y, Chen MT, et al. 2008 Nanosecond electric pulse-induced calcium entry into chromaffin cells. *Bioelectrochemistry* 73: 1–4; Craviso GL, Choe S, Chatterjee P, et al. 2010 Nanosecond electric pulses: a novel stimulus for triggering Ca<sup>2+</sup> influx into chromaffin cells via voltage-gated Ca<sup>2+</sup> channels. *Cell Mol Neurobiol* 30: 1259–1265]. Two such studies also implicate the voltage-gated sodium channels as having roles. These findings show, therefore, that the direct target of the nanosecond pulses is the voltage sensor of these channels and show therefore, that these are not thermal effects. Consequently, it is not surprising that the "safety guidelines" do not predict effects produced from nanosecond pulses, because these "safety guidelines" do not predict non-thermal effects.

When I started to research the issue of effects of nanosecond pulses, this area seemed to be straightforward. Nanosecond pulses produced effects, produced at least in part via VGCC activation, effects that were not predicted by the "safety guidelines." However, when one looks at the nanosecond literature, it is clear that other important types of studies clearly document additional findings that also conflict with "safety guideline" predictions. One of these is that two nanosecond pulses of identical polarities can produce supra-additive effects when the two occur within a few microseconds of each other [Semenov et al. 2018 Electroporation of cells by closely spaced paired nanosecond-range pulses. *Bioelectrochemistry* 121: 135-141]. The second is that when one studies paired nanosecond pulses of *opposite polarities*, the second pulse can greatly depress the effects produced, sometimes called cancellation, by the first pulse [Pakhomov AG, et al. 2014 Cancellation of cellular responses to nanoelectroporation by reversing the stimulus polarity. *Cell Mol Life Sci* 71:4431-4441; Gianulis EC, et al. 2015 Electroporation of mammalian cells by nanosecond electric field oscillations and its inhibition by electric field reversal. *Sci Rep* 2015 Sep 8;5:13818. Doi: 10.1038/srep13818; Sözer EB, Vernier PT. 2019 Modulation of biological responses to 2 ns electrical stimuli by field reversal. *Biochim Biophys Acta Biomembr*. 2019 Apr 11. pii: S0005-2736(19)30077-X. doi: 10.1016/j.bbmem.2019.03.019]. Here again, the pulses must occur within a few microseconds of each other.

"Safety guidelines" do not allow for either supra-additive or depressive effects of a second pulse and do not take into consideration the polarity of exposures and are, therefore, yet again deeply flawed – they falsely assume that all exposures act additively such that average intensities predict effects. This flaw falsifies another assumption of the safety guidelines. When those guidelines assume that EMFs always act additively, one of the underlying assumptions behind that is that EMFs act as scalar variables. What the polarity effects show, however is that EMFs are vector variables, having direction as well as intensity and that polarity of each vector is also important. The important roles of polarity here, shows the vector nature of EMFs and the importance of that vector nature to these paired nanosecond pulse studies. It was known in the 19<sup>th</sup> century that EMFs have vector properties, not scalar properties. It can be seen from this, that the "safety guidelines" are inconsistent with the physics, not just the biology. We have then, different types of findings regarding pairs of nanosecond pulses, each of which is completely inconsistent with predictions of "safety guidelines." 5G is designed to be particularly highly pulsed in order to carry extraordinarily high amounts of information, so that 5G will inevitably have trillions of nanosecond pulses. It follows that using "safety guidelines" to predict effects of 5G radiation is

even more problematic than using “safety guidelines” to predict effects of other types or radiation.

I do not think it is surprising, that pairs of nanosecond pulses that occur within a few microseconds of each other may produce either supra-additive or depressive effects, depending on the polarity of the second pulse. The primary direct target of these pulses is the voltage sensor of the VGCCs and other voltage-gated ion channels, as discussed above such that the properties of the voltage sensor predict how it may be expected to behave in response to EMF exposures. The voltage-sensor has four alpha helices, each designated an S4 helix and with each S4 helix having 5 positive charges, with the 4 S4 helices together making up the voltage sensor. Most of those positive charges are 3 amino acid residues apart from each other, such that the closest charged residues stick out from the helix pretty much on the same side of the helix. Three of those positive charges in each S4 are electrostatically attracted to negative residues on other helices thought to be in fixed positions. What is thought to happen in activation is that there a ratcheting of the S4 helices toward the extracellular space, ratcheting such that the negative charges are now bound to a positive charge 3 residues away from the one that was previously bound. The ratcheting also produces some turning of each S4 helix. This needs to occur several times on each of the four S4 helices to open the channel and allow calcium ions to flow. This ratcheting may occur in response to nanosecond pulse exposures, but the actual secondary structural changes that occur in the voltage-gated ion channel may take much longer than does the ratcheting process. Consequently, the effects of a second pulse, depending on its relative polarity compared with the first pulse, can interact over time periods shorter than the time required for the secondary structural change required to open of the channel.

It should be noted that studies of pulses in the microsecond or millisecond range have not been shown to produce either the supra-additive effects of pulses of identical polarity or the lowered effects produced by a second pulse of opposite polarity. This may be because the relative timing of the two pulses is too far apart.

6. There is also a large literature on the existence of exposure intensity windows where certain specific ranges of intensity of a particular EMF, ***produce maximum biological effects and where ranges either lower or higher produce much lower effects***. The consequences of these findings is that dose response curves are non-linear and are also non-monotone, that is they do not always increase with increasing exposure nor do they always decrease with decreasing exposure. Therefore, the ICNIRP, US FCC, EU and other similar "safety guidelines" are fatally flawed for still an additional reason. I am listing here a series of studies that have reviewed studies of this type. Some of these are genuine review articles and some are primary literature articles that have reviewed substantial amounts of earlier literature. One of the things that is striking here, is that many of these studies have found exposure windows that occur at levels 3, 4 or 5 or more orders of magnitude below the safety guideline cutoffs. Consequently, one can get not just effects but large effects when an exposure window occurs are levels on the order of 100,000 times below allowable "safety guideline" levels. So again, the "safety guidelines" give us absolutely no assurance of safety.

- a. Pall, M. L. 2015 Scientific evidence contradicts findings and assumptions of Canadian Safety Panel 6: microwaves act through voltage-gated calcium channel activation to induce biological impacts at non-thermal levels, supporting a paradigm shift for microwave/lower frequency electromagnetic field action. Rev. Environ. Health 3, 99-116. doi: 10.1515/reveh-2015-0001.
- b. Belyaev, I., 2005. Non-thermal biological effects of microwaves. Microwave Rev. 11, 13-29.
- c. Belyaev, I., 2015. Biophysical mechanisms for nonthermal microwave effects. In: Markov M.S. (Ed), Electromagnetic Fields in Biology and Medicine, CRC Press, New York, pp 49-67.



- d. Adey WR. 1980 Frequency and power windowing in tissue interactions with weak electromagnetic fields. Proc IEEE 68, 119-125.
- e. Blackman CF, Kinney LS, House DE, Joines WT. 1989 Multiple power density windows and their possible origin. Bioelectromagnetics 10:115-128.
- f. Panagopoulos DJ, Margaritis LH. 2009 Biological health effects of mobile telephone radiations. Int J Med Biol Front 15:33-76.
- g. Persson BRR, Eberhardt J, Malmgren L, Persson MB, Brun A, Salford LG. 2005 Effects of microwaves from GSM mobile phones on blood-brain barrier and neurons in rat brain. PERS Online 1:638-641.
- h. Wei Q, Cao ZJ, Bai XT. 2005 [Effect of 900 MHz electromagnetic fields on the expression of the GABA receptor of cerebral cortex cortical neurons in postnatal rats] Wei Sheng Yan Jiu 34: 546-548.
- i. Markov MS. 2004 Myosin light chain modification depending on magnetic fields II. Electromagn Biol Med 23:125-140.
- j. Thompson CJ, Yang YS, Anderson V, Wood AW. 2000 A cooperative model for Ca<sup>++</sup> efflux windowing from cell membranes exposed to electromagnetic radiation. Bioelectromagnetics 21:455-464.

7. Another important factor in determining EMF responses is the type of cell being studied. The relevant studies documenting the importance of cell type are studies where different cell types were studied by the a specific research group using identical methodologies and where the different cell types repeatedly responded differently to the same EMF exposures. I reviewed several studies where such findings were obtained in my 2013 study where single strand breaks in cellular DNA were being measured (Pall ML 2013 Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. J Cell Mol Med 17:958-965. doi: 10.1111/jcmm.12088). I also reviewed several studies of this type when reviewing various genotoxicity studies in my 2015 study (Pall, M. L. 2015 Scientific evidence contradicts findings and assumptions of Canadian Safety Panel 6: microwaves act through voltage-gated calcium channel activation to induce biological impacts at non-thermal levels, supporting a paradigm shift for microwave/lower frequency electromagnetic field action. Rev. Environ. Health 3, 99-116. doi: 10.1515/reveh-2015- 0001). Belyaev IY (2010 Dependence of non-thermal effects of microwaves on physical and biological parameters. Eur J Oncol Library 5: 187-217) reviewed earlier a number of studies, on pp.202 & 203, showing that non-thermal EMF effects were cell type specific. It has repeatedly been found in such studies that stem cells are unusually sensitive to EMF exposures, producing effects where most other cell types do not. Some of these studies have been reviewed by Dr. Belyaev and his colleagues (Belyaev IY, Markovà E, Hillert L, Malmgren LO, Persson BR. 2009 Microwaves from UMTS/GSM mobile phones induce long-lasting inhibition of 53BP1/gamma-H2AX DNA repair foci in human lymphocytes. Bioelectromagnetics 30:129-141. doi: 10.1002/bem.20445; Markovà E, Malmgren LO, Belyaev IY. 2010 Microwaves from Mobile Phones Inhibit 53BP1 Focus Formation in Human Stem Cells More Strongly Than in Differentiated Cells: Possible Mechanistic Link to Cancer Risk. Environ Health Perspect 118:394-399. doi: 10.1289/ehp.0900781). These cell-type specific findings clearly show that that effects are produced via cell type specific biological processes and consequently all claims that are made that one can predict effects just from the physical properties of the EMFs, ***as the ICNIRP, EU, FCC, Canadian and other safety guidelines do, are fraudulent.***

8. The last of these are findings have shown that there are very specific EMF frequencies which produce vastly larger EMF effects than do other slightly different frequencies. These have been interpreted as being due to resonance interactions, where the specific frequency produces a

resonance response in the target involved and therefore produces vastly larger responses. These findings have been reviewed four times, to my knowledge:

- a. Golant MB. 1989 Problem of the resonance action of coherent electromagnetic radiations of the millimeter wave range on living organisms. *Biophysics* 34:370-382. English translation of *Biofizika* 1989; 34:339-348.
- b. Belyaev, I., 2005. Non-thermal biological effects of microwaves. *Microwave Rev.* 11, 13-29.
- c. Belyaev IY. 2010 Dependence of non-thermal effects of microwaves on physical and biological parameters. *Eur J Oncol Library* 5: 187-217.
- d. Belyaev, I., 2015. Biophysical mechanisms for nonthermal microwave effects. In: Markov M.S. (Ed), *Electromagnetic Fields in Biology and Medicine*, CRC Press, New York, pp 49-67.
- e. Adey, WR. 1980 Frequency and power windowing in tissue interactions with weak electromagnetic fields. *Proc IEEE* 68, 119-125.

I would suggest that in animals and plants, the most likely target of such a resonance interaction would be the ion channel voltage sensors. We have no evidence as to whether this is correct or not. Interestingly the only such evidence occurs in the bacterium *Escherichia coli* (reviewed in the second review in the previous paragraph) where the target appears to be the DNA of the cell and where the resonance interaction is influenced by the supercoiling of the DNA.

### Summary of the Evidence

The flaws in the “safety guidelines” can be viewed as falling into four distinct categories: Allowable levels, averaging intensities over a 6 minute or 30 minute period, defects in the "safety guideline" physics and failure of the safety guidelines to take into account the biology. Let's consider those one at a time.

Allowable levels: The findings that 9 different important biological effects each are produced at levels well below “safety guidelines”, show that allowable levels are far too high. The findings that forces placed by EMFs on the voltage sensor that controls the VGCCs and other voltage gated ion channels are approximately 7.2 million times the forces placed on singly electrically charged groups in the aqueous phases of our cells and bodies, suggests that the allowable exposure levels are too high by a factor of approximately 7.2 million. The findings that there are exposure windows producing high level effects that are approximately 5 orders of magnitude below those allowed by the safety guidelines again, suggest a similar estimate to the 7.2 million factor. After all, allowable levels should always be far below levels that can be shown to produce high level effects. These estimates are similar to those proposed in the 2007 Bioinitiative Report. The frequency windows produce effects are many orders of magnitude lower intensities than the While the flaws in setting allowable levels are profound, in terms of the structure of the safety guidelines the assumption that intensities averaged over a 6 minute or 30 minute period can be used to assess safety may well be more profoundly flawed. Let's look at each of 7 relevant findings. 1. There are 13 different reviews each showing that pulsed EMFs produce, in most cases, much greater effects than do non-pulses, also known as continuous wave EMFs of the same average intensity. These findings clearly show that you cannot use average intensities to predict biological effects and therefore safety. But that is, of course, what the “safety guidelines” do. 2. The intensity exposure window studies, showed that there are what have been called intensity windows where a short intensity range of exposures to a particular EMF, produce maximum effects but lower or higher intensities, produce much lower effects. Averaging can only be predictive of biological effects when one has linear dose-response curves. However the window effects studies clearly show that dose-response curves are neither linear nor monotone. They do not always increase with increasing exposure nor do they decrease with decreasing exposure. This clearly shows that exposures averages over 6 minutes or 30 minutes cannot predict biological effects and therefore safety and argues that averaging over any time period

cannot be used in genuine safety guidelines. 3. The nanosecond pulse and paired nanosecond pulse studies show several other flaws behind "safety guideline" averaging. Individual nanosecond pulses repeatedly produce effects but when those intensities are averaged over a 6 minute period, "safety guidelines" predict they cannot produce effects. If you take a typical, let's say 40 nanosecond pulse, averaging its intensity over 6 minutes means you are averaging it over a period approximately  $10^{10}$  times longer than it takes to produce effects, something that makes absolutely no sense. 4-6. Paired nanosecond pulses, where those pulses occur within a few microseconds of each other produce three other failures for averaging. When the second pulse is of the same polarity as the first one, the two together produce supra-additive effects, but these are not predicted by the "safety guidelines." When the second pulse is of the opposite polarity as compared with the first one, the two together produce much lower effects than do the first one alone, such that the second such pulse is said to "cancel" part and often most of the effect of the first pulse. Such cancellation is not, of course predicted from the "safety guidelines" which predict that multiple pulses must act in additive fashion. How, then, should we interpret the failures of the "safety guidelines" to predict either of these two types of paired nanosecond pulse studies? The "safety guidelines" average the intensities of the EMFs as if they were scalars not vectors. Scalars have intensity but no direction whereas EMFs have been known for approximately 200 years to be vectors. Those vectors also have polarity, with the polarity being produced by the angle of the vibrations of the electrical and magnetic fields, perpendicular to the direction of the vector. When the "safety guidelines" average the EMFs as if they were scalars, not vectors, the "safety guidelines" are making a profound error with regard to the physics of the EMFs and therefore the mathematics of averaging. This is not the only profound error in the physics of the "safety guidelines." Another important error concerns the failure of the "safety guidelines" to assess the properties of the voltage sensor, the primary direct target of the EMFs in producing biological effects, including its profound sensitivity to the electrical forces of the low intensity EMFs. 7. The last of these are the frequency windows, where exposures within a frequency window, produce effects at levels many orders of magnitude lower than are needed to produce effects at nearby frequencies. These have been ascribed to resonance interactions with a specific biological target. It can be argued that which it is clear that "safety guidelines" fail to predict these, that because of the high level of specificity of the frequencies involved, that the biological importance may of these failures may have limited importance. That may be true but it may not be true. It has been suggested that the 2.45 GHz frequency commonly used for Wi-Fi is close to a frequency window, such that it may account for the activity of this frequency range. Furthermore, when one has a square wave pulse, and when does a Fourier analysis of such a pulse, one gets an almost infinite number of frequencies making up the square wave, such that some of them may well fall into a frequency window. And it should be noted that digital technologies produce square waves all the time. Dr. Waldmann-Selsam et al (2009. Warning against adverse health effects from the operation of digital broadcast television stations (DVB-T): letter from 3 German physicians to the U.S. President and Congress, <http://www.stayontruth.com/warning-against-adverse-health-effects-digital/TV.php>) wrote a letter stating that digital broadcasting was in their experience, much more damaging to their patients than was the analog technology that it replaced. There may be a similar issue with regard to pulsations, such that the shorter a pulsation, the more complex the Fourier analysis of it will be, such that it is possible that much of the activity of short pulsations may be caused by frequency window effects. The resonance interactions that are the apparent explanation of frequency window effects provide another failure of the "safety guideline" physics. By failing to consider interactions with specific highly sensitive biological targets, "safety guideline" physics is again deeply flawed. Of these "safety guideline" flaws produced by the use of average intensities over a 6 minute or 30 minute period in these "safety guidelines," the first six of these flaws discussed here should be viewed as fatal flaws because they produce large errors in assessment of very common exposures. The 7<sup>th</sup>,

the frequency window flaw, its uncertain importance in producing failed predictions of the "safety guidelines" means that it is unclear whether it should be considered a fatal flaw or not. In summary, using this same definition of fatal flaws, we have two fatal flaws in the "safety guideline" physics, three fatal flaws in the safety guideline allowable levels, six fatal flaws in using average intensities over a 6 minute or 30 minute period. Each of these 11 fatal flaws and the additional flaws documented from frequency window studies, because each of them is documented through the biology, show that no valid safety guidelines can be produced without paying close attention to the biology, but that is exactly what they current "safety guidelines" do. There is, an additional finding which emphasizes that the biology cannot be ignored. That additional finding is that when particular research groups look at multiple cell types using the same methodology, they often find that different cell types respond very differently to a particular EMF exposure. Consequently the heterogeneity of different cell types cannot be ignored.

The collective failures of the "safety guidelines" documented here, may well be the most massive failure in the history of science to pay attention to the science. They may well be the most massively consequential such collective failure, because of the vast proliferation of technologies based on these safety guidelines in almost every country on earth, consequential because of both the human impact and ecological impacts of the EMF effects. Each of these eight distinct types of findings show that the ***safety guidelines are fraudulent because they do not predict biological effects***. While this document is aimed at the 1998 ICNIRP, US FCC, EU, Canadian and Australian and US FCC safety guidelines, the similar safety guidelines produced by other countries, including 2013 UK guidelines are similarly fraudulent for each of those same eight reasons. ***All guarantees of safety given by these organizations or by industry organizations, which are based on these or similar safety guidelines, are similarly fraudulent.*** The fraudulence is caused, in part, by the series of false assumption underlying these safety guidelines: Assumptions that average intensities or average SAR can be used to assess safety are false. Assumptions that one can ignore pulsations including very short spikes and nanosecond pulses are false. Assumptions that you one ignore biological heterogeneity and assess effects simply based on physics are false. Assumptions that dose-response curves are linear or at least monotone are false. Assumptions that there are no mechanisms that can explain the existence of non-thermal effects are false. Assumptions that electrical forces produced by low intensity EMFs are too weak to do anything are false. Assumptions that identifying average SARs in a specific study is of special importance in judging the quality or biological relevance of the study are false. The consequences of all this, is that we have a multi-trillion dollar (or multi-trillion euro) set of industries, the telecommunication industries where all assurances of safety are based entirely on massive fraud.

I wish that there were some simple modification of these or other guidelines that could provide more reliable safety guidelines, but no such modification exists. At this point in time, the only way to determine biological safety is to do biological safety testing. And biological safety testing is very challenging because of the important roles of:

1. Complex dose-response curves, with regard to intensities and frequencies. The exposure windows create particular challenges, because of high level effects produced by levels of exposure as much as 5 orders of magnitude below current safety guidelines.
2. Biological heterogeneity is important as shown by different responses of different cell types.
3. The physics is complex both with regard to the vector nature of EMFs and the stunning sensitivity of the main direct target of EMFs, the voltage sensor, to the electrical forces of weak EMFs.

4. The complex roles of pulsation as shown by studies of nanosecond, microsecond and millisecond pulses.

### What About 5G?

5G will entail using millimeter wave EMFs with sufficient band width and extraordinary levels of pulsation to wirelessly communicate many orders of magnitude greater amounts of information per unit time than do current wireless communication systems. This is clearly stated in the paper that I think of as a propaganda document which, then, fails to take into account any of the established findings previously discussed in this document (Wu T, Rapaport TS, Collins CM. 2015 Safe for generations to come. IEEE Microw Mag. 2015 March ; 16(2): 65–84.) The extraordinary 5G health problems are created, in part, by the extraordinary pulsation levels. We not only have the 13 reviews each of which show that pulsed EMFs are, in most cases much more biologically active than are non-pulsed EMFs of the same average intensity but we also have the nanosecond pulse studies that are particularly relevant to 5G. The nanosecond pulse studies, described above are particularly relevant to 5G because of the extraordinary pulsation levels of 5G antennae communicating with the “internet of things” will inevitably involve astronomical numbers of nanosecond pulses. The nanosecond pulses are relevant because the individual nanosecond pulses produce effects not predicted by the “safety guidelines.” They are also especially relevant because of the supra-additive effects seen when nanosecond pulses of identical polarity occur within a few microseconds of each other and these supra-additive effects are also not predicted by the “safety guidelines.”

Additional 5G health problems are also created in two ways, by the millimeter wave frequencies. The electrical parts of millimeter waves are highly absorbed by materials including the materials in our buildings and the materials in our bodies. Such absorption means that the 5G plans have entailed putting out tens of millions of antennae in close proximity to our homes and other buildings, putting out very high power EMFs, such that the electrical parts can penetrate into our homes and other buildings such that electrical devices can communicate with the 5G antennae. The penetration into buildings will entail use of phase arrays which produce an additional type of pulsation to which we are to be exposed. The high absorption argues that these millimeter waves will be particularly active in activating the VGCCs because the mechanism of such absorption involves interacting with electrically charged groups, including the electrically charged groups of the voltage sensor. That is one very large problem, very high level VGCC activation. The second large problem is the high numbers and high power of the so called small cell antennae which means that it will be essentially impossible to avoid the 5G EMFs, especially when we are outside and have no shielding between our bodies and the antennae. I predict, therefore, that 5G will inevitably create not only human but also ecological disasters of unparalleled proportions. Small mammals and birds and insects will be heavily impacted because of their large surface to volume ratios. For example, two US Patents # 6,073,365 and # 6192598 each showed that millimeter waves have potent insecticidal activity and that the higher the frequency, the higher the insecticidal activity. Plants are also highly susceptible because even large trees have their leaves and reproductive organs highly exposed. One of the consequences that I predict is that we will have huge conflagrations because EMFs make plants vastly more flammable. That may make fires much worse than the recent California fires commonplace. I am, therefore, profoundly concerned about both the human effects and the ecological effects.

Very high level surface effects in humans will, of course, also occur and will probably include:

1. High incidences of melanomas, given multiple reports that EMFs cause melanomas in humans, including in parts of the body not commonly exposed to sunlight. Other

- expected cancers would also include leukemias and lymphomas because the precursor cells circulate in the blood up through blood vessels near the surface of the body.
2. Erythrocytes are impacted by EMFs, forming such aberrant structures as rouleau bodies, spike structures on the surface of the erythrocytes and hemolysis. Hemolysis will produce anemia. Any and all of these may be expected from 5G exposures.
  3. Electromagnetic hypersensitivity often involves surface reddening and pain from EMF exposures, so these are likely to be very common consequences of 5G.
  4. Autoimmune diseases can also be produced by EMFs, acting via increased T-cell calcium signaling to produce autoimmunity. Again because the T-cells circulate near the surface of the body, large increases in autoimmunity may be predicted as a consequence of 5G radiation.

One of the claims that the industry makes is that millimeter wave frequencies to be used in most 5G radiation will be absorbed in the outer 1 mm of the body and that therefore, 5G will have no effects in underlying tissues. I showed in Chapter 7 of my 90 page document what the industry claims about penetration of microwave effects were false such that microwaves act at least 40 times more deeply in the body than the industry claims is possible. The way the physics can produce such deeply penetrating effects was also discussed. I predicted therefore that millimeter waves will act deeply as well. Now we have evidence from two CIA translated documents that millimeter waves act at least 20 times more deeply in the body than the industry claims is possible (Zalyobokskaya NP, 1977. Biological effect of millimeter radiowaves. Vrachebnoye Delo 3: 116-119. Declassified and Approved for release 2012/05/10: CIA-RDP88B01125R000300120005-6; Levedeva NN, Reactions of the central nervous system to peripheral effects of low-intensity EHF emissions. Approved for release 2000/08/10: CIA-RDP96-00792R000100070001-9).

The first of these documents shows that internal organs of rodents including heart, kidney, liver spleen and bone marrow can be heavily impacted by low intensity continuous millimeter wave radiation. There are even more severe effects on the skin, as may be expected. These studies show that these millimeter wave EMFs produce effects at least 20 times deeper than the industry claims is possible. The effects seen start out as modest effects that can be reversible with cessation of exposure but become much more severe with increasing times of exposure. The human study (second paper) was an EEG study where electrical activity in the brain was being monitored. Here for the low intensity millimeter continuous wave EMF exposure to have effects, it must penetrate the hair, skin, skull and meninges surrounding the brain. Again, in humans effects are found at least 20 times deeper than the industry claims is possible. Cardiac effects of millimeter waves, discussed below, also show much deeper penetration of effects of such waves than predicted by industry.

How then are these highly penetrating effects produced? Although the electrical parts of the EMFs may be absorbed readily, the magnetic parts are very highly penetrating. These can then put forces on dissolved ions in the aqueous phases of our cells and tissues, moving them and regenerating the electrical parts of the EMFs, with the same frequency and same pulsations, just with much lower intensity. However with the voltage sensor of the VGCCs so stunningly sensitive to the electrical forces, this can produce effects very deeply within the body.

These deep effects produced by the millimeter wave EMFs in these two CIA documents, no doubt deeply underestimate that genuine 5G radiation will produce, given that 5G produces extraordinarily high level of pulsations. What effects do I predict for genuine 5G? I predict that similar but much more severe effects will be produced by 5G as are produced by microwave frequency EMFs. I also predict because of the roles of aqueous dissolved ions in producing these

deep effects, that regions of the body with large such internal "bodies of water" may be expected to produce particularly severe problems. These may include:

- a) Various types of birth defects because of the role of the amniotic fluids and the increased extracellular water content in the tissues of the fetus.
- b) Massive epidemics of blindness due to the role of the aqueous and vitreous humors of the eye.
- c) Massive epidemics of kidney failures due to the water in the kidney.
- d) Larger epidemics of life threatening cardiac changes in the electrical control of the heart, because of the large blood fluids in the heart.
- e) Large epidemics of circulatory problems, possibly including aortic and other arterial aneurisms.

The only way to test for these and other health impacts is to do biological testing with genuine 5G radiation with all of the pulsations that will be involved once it is connected to the "internet of things." Those many billions of connections imply astronomical numbers of pulsations including nanosecond pulsations. Biological testing of genuine highly pulsed 5G radiation is exactly what the regulatory agencies and the industry are both avoiding doing.

There is some evidence with regard to d, above, from worrisome cardiac effects produced by non-pulsed millimeter wave frequency radiation on animals. Chernyakov GM, Korochkin VL, Babenko AP, Bigdai EV. 1989 Reactions of biological systems of various complexity to the action of low-level EHF radiation. In Devyakov ND, (ed.) Millimeter Waves in Medicine and Biology. Moscow: Radioelectronica pp. 141-167. (in Russian) found that millimeter waves could produce tachycardia and arrhythmia in the hearts of frogs, even when the hearts were detached from neuronal control. Potekhina IL, Akoyev GN, Yenin LD, Oleyner (1992 Effects of low-intensity electromagnetic radiation in the millimeter range on the cardiovascular system of the white rat. Fiziol Zh 78:35-41 (in Russian)) found that 20 minute exposures to non-thermal non-pulsed millimeter wave EMFs of certain frequencies caused pronounced arrhythmia in the rat. 3 hour exposures caused about 25% of the animals to undergo apparent sudden cardiac death. You will note that earlier in this paper, the possible causation of sudden cardiac death in humans from EMF exposures was discussed. These two studies of millimeter wave cardiac effects also showed, as did other studies discussed above, that the millimeter waves can act vastly more deeply in the body than the industry claims is possible.

These animal studies on non-pulsed millimeter wave effects, discussed in the previous paragraph, make very plausible some animal effects that were reported to occur in The Netherlands during a recent introduction of 5G (Massive starling death by 5G? #FactCheck. <http://wearechange.nl/?p=729>). Here starling birds died suddenly over a several day period of possible sudden cardiac death during the approximate time period of 5G introduction. This is still a matter of controversy and I am not making any conclusions here. But what is clear is that the finding that non-pulsed millimeter wave frequency EMF exposures can cause cardiac effects in animals including apparent sudden cardiac death makes it much more plausible that similar observations in birds may have been caused by 5G radiation during 5G testing.

I am simply going to repeat a statement I made earlier. The "safety guidelines" have been known to be bogus for over 40 years, based on findings of non-thermal effects at levels of exposure well below those allowed under our "safety guidelines." What is clear from this document, is that every single aspect of the safety guidelines is fatally flawed for multiple reasons. "Safety guidelines" using averaged intensities and SAR (or average power densities, as some recent proposals suggest) do not predict biological effects and therefore do not predict safety for over 20 distinct reasons. There are multiple fatal flaws that destroy the "safety guideline" allowable

levels, the use of average intensities over 6 minutes or 30 minutes and the "safety guideline" physics, flaws that are produced by the complete failure to pay any attention to any of the biological data that can be used to test the predictions of the safety guidelines. Putting out tens of millions of 5G antennae, in close proximity to almost every home or other building, making it impossible to avoid massive exposures, based on fraudulent "safety guidelines" without doing one single biological test of 5G radiation with all of the pulsations it will entail, is at a minimum, a candidate for being the stupidest thing that has ever been done.